




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07D 257/02, 493/10, 405/14, A61K 49/00, C07F 9/6524 // (C07D 493/10, 317:00, 317:00)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/31444</b> <b>(43) International Publication Date:</b> 23 November 1995 (23.11.95)
<b>(21) International Application Number:</b> PCT/IB95/00337 <b>(22) International Filing Date:</b> 9 May 1995 (09.05.95)  <b>(30) Priority Data:</b> 08/241,253      11 May 1994 (11.05.94)      US  <b>(71) Applicant:</b> BRACCO INTERNATIONAL B.V. [NL/NL]; De Boelelaan 7, NL-1083 HJ Amsterdam (NL).  <b>(72) Inventors:</b> RANGANATHAN, Ramachandran, S.; 196 Sayre Drive, Princeton, NJ 08648 (US). PILLAI, Radhakrishna; 39 Donald Avenue, Kendall Park, NJ 08826 (US). RATSEP, Peter, C.; 24 Mario Drive, Hamilton Square, NJ 08690 (US). SHULKA, Rajesh; 23 Fairway Court, Lawrenceville, NJ 08648 (US). TWEEDLE, Michael, F.; 72 Library Place, Princeton, NJ 08540 (US). ZHANG, Xun; 4 Stanford Drive, Kendall Park, NJ 08824 (US).  <b>(74) Agent:</b> TOMOV-GRISOGONO, Nenad; Bracco Research S.A., 7, route de Drize, CH-1227 Carouge (CH).		<b>(81) Designated States:</b> CA, CN, CZ, JP, KR, NO, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ENHANCED RELAXIVITY MONOMERIC AND MULTIMERIC COMPOUNDS  <b>(57) Abstract</b> <p>Metal chelates capable of exhibiting an immobilized relaxivity between about 60 and 200 mM<sup>-1</sup>s<sup>-1</sup>/metal atom are useful as magnetic resonance imaging agents. Additionally, a compound which is useful as a metal-chelating ligand has formula (II): wherein Q is a 4- to an 8-membered carbocyclic ring which may be fully or partially saturated; t is an integer from 2 to 16; each R group is independently hydrogen, -OH, -CH<sub>2</sub>-A, -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A or a functional group capable of forming a conjugate with a biomolecule, provided that at least two of the R groups are selected from -CH<sub>2</sub>-A or -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A; and A is a moiety capable of chelating a metal atom.</p> <div style="text-align: right; margin-top: 20px;">  <p style="margin-left: 10px;">(II)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -


ENHANCED RELAXIVITY MONOMERIC AND MULTIMERIC  
COMPOUNDS

Brief Description of the Invention

5 In accordance with the present invention, novel monomeric and multimeric compounds having enhanced relaxivities are provided. These compounds are useful, for example, as metal-chelating ligands. The compounds are also useful in the form of metal  
10 complexes as diagnostic contrast agents. When the metal in the complex is paramagnetic, the diagnostic contrast agents are especially suitable for magnetic resonance imaging (MRI).

In one embodiment of the invention, certain  
15 specific compounds comprise a tetraazacyclododecane macrocycle, and are represented by the formula I:

I



20 wherein

each m, n, o and p is independently 1 or 2;

- 2 -

- q is 0 or 1;  
each G is independently -COOR'', -P(O)(OR'')<sub>2</sub>,  
-P(O)(OR'')(R'') or -C(O)N(R'')<sub>2</sub>;  
each R' is independently hydrogen or alkyl,
- 5 alkoxy, cycloalkyl, hydroxyalkyl or aryl, each of  
which is optionally substituted, or a functional  
group capable of forming a conjugate with a  
biomolecule or of forming a multimer of said compound  
of formula I;
- 10 each R'' is hydrogen;  
each R<sup>13</sup> through R<sup>20</sup> is independently  
hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl or a  
functional group capable of forming a conjugate with  
a biomolecule or of forming a multimer of said  
15 compound of the formula I;  
or R<sup>13</sup> together with R<sup>15</sup>, and R<sup>17</sup> together with  
R<sup>18</sup>, independently form, together with the carbon  
atoms in the tetraazacyclododecane macrocycle to  
which they are attached, a fused fully or partially  
20 saturated non-aromatic cyclohexyl ring which may be  
unsubstituted or substituted by one or more halogen,  
alkyl, ether, hydroxy or hydroxyalkyl groups, and  
which may be further fused to a carbocyclic ring, or  
R<sup>13</sup> and R<sup>15</sup> are each hydrogen and R<sup>17</sup>, together with  
25 R<sup>18</sup>, forms a fused fully or partially saturated non-  
aromatic cyclohexyl ring as defined above, or R<sup>13</sup>,  
together with R<sup>15</sup>, forms a fused fully or partially  
saturated non-aromatic cyclohexyl ring as defined  
above and R<sup>17</sup> and R<sup>18</sup> are hydrogen; provided that  
30 (a.) when G is always -COOR'' and (i.) R', R'', R<sup>14</sup>  
and R<sup>16</sup> through R<sup>20</sup> are all hydrogen, then R<sup>13</sup> and R<sup>15</sup>  
are other than hydrogen; (ii.) R'' and R<sup>13</sup> through R<sup>20</sup>  
are all hydrogen, and m, n, o, p and q are each 1,  
then (CR'R') is other than (CH<sub>2</sub>) and (CHCH<sub>3</sub>); (iii.)

- 3 -

R', R'', R<sup>13</sup>, R<sup>14</sup>, R<sup>17</sup> and R<sup>20</sup> are all hydrogen, then at least two of R<sup>15</sup>, R<sup>16</sup>, R<sup>18</sup> and R<sup>19</sup> are other than methyl; and (iv.) R'', R<sup>16</sup>, R<sup>19</sup> and R<sup>20</sup> are all hydrogen, and each (CR'R') is independently (CHR') or (CH<sub>2</sub>CHR'), then R<sup>13</sup> and R<sup>15</sup>, and R<sup>17</sup> and R<sup>18</sup>, are other than a fused ring; and (b.) when G is always -P(O)(OR'')<sub>2</sub>, -P(O)(OR'')(R'') or -C(O)N(R'')<sub>2</sub>, then at least one R' or R<sup>13</sup> through R<sup>20</sup> is other than hydrogen;

10 or a salt or multimeric form thereof.

#### Detailed Description of the Invention

Listed below are definitions of various terms used in the description of this invention. These definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific instances, either individually or as part of a larger group.

The expression "relaxivity" refers to the effectiveness of a metal chelate to reduce the relaxation time of bulk water in contact with the metal chelate.

The expression "immobilized relaxivity" refers to the relaxivity measured when a chelate moiety can undergo only slow molecular reorientation because of rigid attachment to a large moiety or a physiological surface, or because it is dissolved in a medium of high viscosity.

The expression "water relaxivity" refers to relaxivity in water where a chelate moiety possesses a relaxivity dominated by overall molecular reorientation.

The expression "enhanced relaxivity" refers to relaxivity values made greater than those of well

- 4 -

characterized prior art molecules by 1.) altering the electronic relaxation rate,  $\tau_s$ , through modifications of the metal-donor atom bond vibration frequencies and/or amplitudes, (this being  
5 accomplished, for example, by increasing the steric bulk and/or orientation of organic elements bonded to the donor atoms and/or the macrocyclic carbon atoms),  
2.) in a multimer by decreasing the internal molecular motion of one monomer unit relative to  
10 another (this being accomplished, for example, by increasing the steric bulk of the organic groups linking the monomer units) or 3.) by decreasing the molecular reorientation of a monomer or a multimer attached to a large moiety or a physiological  
15 surface.

The term "stability" refers to the equilibrium formation constant ( $K$ ) of the reaction  $M+L \rightarrow M(L)$  where  $K=[M(L)]/[M][L]$ ,  $M$  is a metal ion,  $L$  is a chelating ligand and  $M(L)$  is a chelate complex of a  
20 metal and a ligand.

The term "alkyl" refers to both straight and branched, unsubstituted chains of carbon atoms. Those chains having 1 to 5 carbon atoms are preferred. Methyl is the most preferred alkyl group.

25 The term "cycloalkyl" refers to cyclic hydrocarbon groups of 3 to 8 carbon atoms. The groups may be unsubstituted or substituted by, for example, alkyl, halogen, hydroxy, hydroxyalkyl, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthiol,  
30 nitro, cyano, carboxy, carbamoyl, alkoxycarbonyl, alkylsulfonyl, sulfonamido and the like.

The term "alkoxy" refers to  $-\text{alkyl}(O)$ . Methoxy is the most preferred alkoxy group.

- 5 -

The term "aryl" refers to phenyl, pyridyl, furanyl, thiophenyl, pyrrolyl, imidazolyl and the like, all of which may be substituted. Preferred substituted aryl groups are those substituted with 1, 2 or 3 halogen, nitroamino, maleimido, isothiocyanato, hydroxy, hydroxyalkyl, alkyl, alkoxy, carbamoyl, carboxamide, acylamino or carboxy moieties.

"Hydroxyalkyl" refers to straight and branched alkyl groups including one or more hydroxy radicals such as  $-\text{CH}_2\text{CH}_2\text{OH}$ ,  $-\text{CH}_2\text{CH}_2\text{OHCH}_2\text{OH}$ ,  $-\text{CH}(\text{CH}_2\text{OH})_2$  and the like. (See, for example, Sovak, M., Editor, Radiocontrast Agents, Springer-Verlag, 1984, pp. 1-125).

The term "aralkyl" refers to an aryl group bonded through an alkyl group.

The term "carbocyclic ring" refers to a ring system in which all the ring atoms are carbon, e.g., phenyl or cyclohexyl. The ring may be unsubstituted or substituted by, for example, alkyl, halogen, hydroxy, hydroxyalkyl, alkoxy, alkanoyl, alkanoyloxy, amino, alkylamino, dialkylamino, alkanoylamino, thiol, alkylthiol, nitro, cyano, carboxy, carbamoyl, alkoxy carbonyl, alkylsulfonyl, sulfonamido and the like.

The term "halogen" refers to bromo, chloro, fluoro or iodo.

The term "alkanoyl" refers to the group  $\text{alkyl-C(O)-}$ .

The term "alkanoyloxy" refers to the group  $\text{alkyl-C(O)-O-}$ .

The term "amino" refers to the group  $-\text{NH}_2$ .

The term "alkylamino" refers to the group  $-\text{NHR}$  where R is alkyl.

- 6 -

The term "dialkylamino" refers to the group -NRR' where R and R' are each, independently, alkyl.

The term "alkanoylamino" refers to the group alkyl-C(O)-NH-.

5 The term "thiol" refers to the group -SH.

The term "alkylthiol" refers to the group -SR where R is alkyl.

The term "nitro" refers to the group -NO<sub>2</sub>.

The term "cyano" refers to the group -CN.

10 The term "carboxy" refers to the group -C(O)OH or the group -C(O)OR where R is alkyl.

The term "alkoxycarbonyl" refers to the group alkoxy-C-(O)-.

15 The term "alkylsulfonyl" refers to the group alkyl-SO<sub>2</sub>-.

The term "sulfonamido" refers to the group -SO<sub>2</sub>NH<sub>2</sub>, the group -SO<sub>2</sub>NHR or the group -SO<sub>2</sub>NRR' where R and R' are each, independently, alkyl, cycloalkyl or aryl.

20 The term "carbamoyl" refers to the group -C(O)NH<sub>2</sub>, the group -C(O)NHR or the group -C(O)NRR' where R and R' are each, independently, alkyl, alkoxy or hydroxyalkyl.

25 The term "carboxamide" refers to the group -C(O)NH<sub>2</sub>, the group -C(O)NHR or the group -C(O)NRR' where R and R' are each, independently, alkyl.

The term "acylamino" refers to the group -NH-C(O)-R where R is alkyl.

30 The expressions "bioactive group" and "bioactive moiety" denote a group which is capable of functioning as a metabolic substrate, catalyst or inhibitor, or is capable of being preferentially taken up at a selected site of a subject, such as by




- 7 -

possessing an affinity for a cellular recognition site.

When compounds of the formula I are in the multimeric form, each monomer is preferably linked by  
5 a cyclic bridging group represented by the general formula II:

- 8 -

II



wherein


- Q is a 4- to an 8-membered carbocyclic ring  
 5 which may be fully or partially saturated;  
 t is an integer from 2 to 16;  
 each R group is independently hydrogen, -OH,  
 -CH<sub>2</sub>-A, -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A or a functional group  
 capable of forming a conjugate with a biomolecule,  
 10 provided that at least two of the R groups are  
 selected from -CH<sub>2</sub>-A or -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A; and

A is the monomer of formula I.

- Of course, the cyclic bridging group may be used to  
 link known moieties as well. Where it is a known  
 15 moiety which is being linked by the cyclic bridging  
 group of formula II, A is any moiety capable of  
 chelating a metal atom.


Preferred bridging groups of the formula II  
 are those wherein

20



is a compound of the following formula III:

- 9 -



wherein


each  $R^1$  through  $R^{12}$  group is independently hydrogen, -OH, -CH<sub>2</sub>-A, -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A or a

5 functional group capable of forming a conjugate with a biomolecule;

at least two of  $R^1$  through  $R^{12}$  are selected from -CH<sub>2</sub>-A or -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A;

10  $R^8$  and  $R^9$  taken together may additionally form the group -O-[C(RR)]-O- where each R is independently hydrogen or alkyl, or  $R^8$  and  $R^9$  taken together may form


- 10 -



and

A is a moiety described above.

- 5 When A is a known monomer, for example, A is preferably



wherein

- 10 each R' is independently hydrogen, alkyl, alkoxy, hydroxyalkyl, aryl, aralkyl or arylalkoxy;  
each R'' is hydrogen; and  
each n is 1 or 2;

- 11 -

or a salt thereof.

Contrast agents with significantly enhanced relaxivities are of great interest, not only because they offer improved efficacy at reduced doses for  
5 current clinical applications, but also because they may provide the sensitivities needed for imaging various biochemical processes.

Certain preferred compounds having enhanced relaxivities are (i.) chelates possessing a stability  
10 greater than or equal to  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 60 and  $200 \text{ mM}^{-1}\text{s}^{-1}$ /metal atom, for example between about 70 and  $150 \text{ mM}^{-1}\text{s}^{-1}$ /metal atom, or between about 80 and  $100 \text{ mM}^{-1}\text{s}^{-1}$ /metal atom; and (ii.) multimeric chelates  
15 possessing monomer units with a stability greater than or equal to  $10^{15} \text{ M}^{-1}$ , and having relaxivity values (not immobilized) greater than  $5 \text{ mM}^{-1}\text{s}^{-1}$ /metal atom.

In designing new chelates with these elevated  
20 relaxivities, the inventors have noted that immobilized relaxivity can depend strongly, for example, on the structure of the chelate. Without being bound by any particular theory, apparently a mechanism that involves rigidifying the chelate  
25 structures in solution with, for example, alkyl substitutions on the tetraaza ring of chelates, especially when the substitution is introduced into the carboxylate arms of the chelates, affects immobilized relaxivity. In the Solomon-Bloembergen-  
30 Morgan (SBM) model, the electronic relaxation of a paramagnetic metal complex is viewed as occurring through a dynamic modulation process about the transient zero-field splitting (ZFS) of the metal's electronic spin levels. This transient ZFS is

- 12 -

induced by the structural distortion of the metal complex from its ideal symmetry in solution which is thought to be caused by its collision with solvent molecules. Alkyl substitution on either the tetraaza ring or the carboxylate arms of a chelate is thought to reduce the flexibility of the chelate for structural distortion in solution. This in turn reduces the magnitude of ZFS, giving an increased immobilized relaxivity value.

Water relaxivity is generally determined in aqueous Bis Tris buffer (pH 7) solutions by the standard inversion-recovery method (known to those in the art) at 20 MHz and 40°C. (See, e.g., X. Xhang, Inorganic Chemistry, 31, 1992, 5597, the entire contents of which are hereby incorporated by reference.) While a mathematical description of the relaxation mechanism at the presence of a paramagnetic species is provided by the classical SBM equations, it is experimentally difficult to explore the dependence of relaxivity on structure because the relaxivity values of most low-molecular weight chelates are often controlled by their rapid tumbling motions in regular aqueous solutions.

To eliminate the overriding effect of molecular tumbling in regular aqueous solutions, the relaxivity of a chelate can be determined in aqueous sucrose solutions or other media that result in a reduction of molecular reorientation of any solute. In these solutions, the relaxivity values approximate, under optimal conditions, the relaxivities of chelates in biologically immobilized systems such as those covalently attached to cell surfaces. Hence, these values in sucrose solutions are defined as immobilized relaxivity. Immobilized

- 13 -

relaxivity is generally determined under the same conditions as water relaxivity (see supra) except that the viscosity of the solutions is increased to 80 cp by the addition of solid sucrose to the aqueous solution of chelate, and the temperature is set at 20°C. In the calculation of relaxivity, one uses the concentration of the solute in the aqueous solution (before sucrose is added).

When used for relative comparison, this method serves as a simple screening technique for differentiating chelates designed for efficient biological targeting.

Thus, the compounds of the invention, including compounds of the formula I and formula II, and salts thereof, may be complexed with a paramagnetic metal atom and used as relaxation enhancement agents for magnetic resonance imaging. These agents, when administered to a mammalian host (e.g., a human) distribute in various concentrations to different tissues, and catalyze relaxation of protons (in the tissues) that have been excited by the absorption of radiofrequency energy from a magnetic resonance imager. This acceleration of the rate of relaxation of the excited protons provides for an image of different contrast when the host is scanned with a magnetic resonance imager. The magnetic resonance imager is used to record images at various times, generally either before and after administration of the agents, or after administration only, and the differences in the images created by the agents' presence in tissues are used in diagnosis. In proton magnetic resonance imaging, paramagnetic metal atoms such as gadolinium(III), and manganese(II), chromium(III) and iron(III) (all are paramagnetic metal atoms with

- 14 -

favorable electronic properties) are preferred as metals complexed by the ligands of the invention, including the ligands of formula I and formula II. Gadolinium(III) is the most preferred complexed metal  
5 due to the fact that it has the highest paramagnetism, it has low toxicity when complexed to a suitable ligand, and it has high lability of coordinated water. When the distance between the monomeric gadolinium chelate units in a complex is at least about 6  
10 angstroms, the complexes tend to be sufficiently stable. Those compounds of formula II, when complexed with gadolinium (III) ions, are particularly useful. The distance between the monomeric gadolinium chelate units in these complexes is generally greater than 6  
15 angstroms (although in certain circumstances a distance of 4.5 angstroms is sufficient), and the rigid bridges of these complexes assist in reducing independent motion of the gadolinium ions.

The metal-chelating ligands of the present  
20 invention can also be complexed with a lanthanide (atomic number 58 to 71) and used as chemical shift or magnetic susceptibility agents in magnetic resonance imaging or in magnetic resonance in vivo spectroscopy.

25 While the above-described uses for the metal-chelating ligands of the present invention are preferred, those working in the diagnostic arts will appreciate that the ligands can also be complexed with the appropriate metals and used as contrast  
30 agents in other imaging techniques such as x-ray imaging, radionuclide imaging and ultrasound imaging, and in radiotherapy.



- 15 -

Use in Imaging

To use the ligands of the present invention for imaging, they are first complexed with an appropriate metal. This may be accomplished by methodology known in the art. For example, the metal can be added to water in the form of an oxide or in the form of a halide or acetate and treated with an equimolar amount of a ligand of the present invention. The ligand can be added as an aqueous solution or suspension. Dilute acid or base can be added (where appropriate) to maintain a suitable pH. Heating at temperatures as high as 100°C for periods of up to 24 hours or more may sometimes be employed to facilitate complexation, depending on the metal and the chelator, and their concentrations.

Pharmaceutically acceptable salts of the metal complexes of the ligands of this invention are also useful as imaging agents. They can be prepared by using a base (e.g., an alkali metal hydroxide, meglumine, arginine or lysine) to neutralize the above-prepared metal complexes while they are still in solution. Some of the metal complexes are formally uncharged and do not need cations as counterions. Such neutral complexes may be preferred in some situations as intravenously administered x-ray and NMR imaging agents over charged complexes because they may provide solutions of greater physiologic tolerance due to their lower osmolality.

The present invention also provides pharmaceutical compositions comprising a compound of the invention, including a compound of the formula I or II, or a salt of one of these compounds, optionally complexed with a metal, and a pharmaceutically acceptable vehicle or diluent. The

- 16 -


present invention further provides a method for diagnostic imaging comprising the steps of administering to a host a compound of the invention, or a salt thereof, which is complexed with a metal, and obtaining a diagnostic image, preferably a magnetic resonance image, of said host.

Sterile aqueous solutions of the chelate complexes of the present invention are preferably administered to mammals (e.g., humans) orally, intrathecally and, especially, intravenously in concentrations of about 0.003 to 1.0 molar. The metal complexes of the present invention may be employed for visualization of various sites. For example, for the visualization of brain lesions using magnetic resonance imaging, a gadolinium complex of a ligand of the invention, including a ligand of the formula I or formula II, may be administered intravenously at a dose of 0.001 to 0.5 millimoles of the complex per kilogram of body weight, preferably at a dose of 0.001 to 0.3 millimoles/kilogram. For visualization of the kidneys, the dose is preferably 0.05 to 0.20 millimoles/kilogram. For visualization of the heart, the dose is preferably 0.001 to 0.3 millimoles/kilogram. For visualization of the liver, the dose is preferably 0.001 to 0.3 millimole/kilogram.

The pH of the formulation of the present metal complexes is preferably between about 6.0 and 8.0, most preferably between about 6.5 and 7.5. Physiologically acceptable buffers (e.g., tris(hydroxymethyl)-aminomethane) and other physiologically acceptable additives (e.g., stabilizers such as parabens) may also be present.

- 17 -

It is also advantageous to employ dual scavenging excipients such as those described in copending application U.S. Serial No. 032,763, filed March 15, 1993, entitled "DUAL FUNCTIONING EXCIPIENT FOR METAL CHELATE CONTRAST AGENTS", incorporated herein by reference. Those excipients have a general formula corresponding to:



wherein D and D' are independently Ca or Zn, L' is an organic ligand which may be different from, or the same as, the ligand employed to complex the metal, and s and t are independently 1, 2 or 3.

As already noted, the present invention further includes multimeric forms of the compounds of the invention, including those of formula I and formula II, such as dimers, trimers, tetramers, etc. Known functional groups and technology are readily useable to provide such multimers.

Compounds of the present invention may include those containing functional group(s) capable of forming a conjugate with a biomolecule. These compounds are preferably chelates, including a functional group, capable of exhibiting an immobilized relaxivity between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom, for example between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom, or between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom. Similarly, the chelates, once conjugated to a biomolecule of size greater than or equal to about 40,000 daltons, are also preferably capable of exhibiting a relaxivity between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom, for example between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom, or between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom. Preferred biomolecules are

- 18 -

peptides, polypeptides and oligosaccharides or fragments thereof, although other biomolecules such as proteins, particularly monoclonal antibodies, lipids, sugars, alcohols, bile acids, fatty acids, receptor-binding ligands, amino acids and RNA, DNA or modified fragments of these may be conjugated to the compounds of the present invention. For smaller biomolecules, the enhanced relaxivity afforded by the chelates of this invention may be more fully realized when the chelate-biomolecule conjugate becomes immobilized *in vivo*, such as by binding to a receptor on a cell surface or by binding to another biomolecule.

Conjugates where a compound of the invention, or salt and/or multimer thereof, is linked to a biomolecule such as a protein, provided by the present invention, are novel, as are metal complexes and pharmaceutical compositions containing, and methods of using (e.g., for imaging), the aforementioned conjugates. Conjugation may be achieved *in vitro*, such as by use of known conjugation methodologies, or *in situ* in a living subject by administration of a compound containing one or more of the aforementioned functional groups.

For linking the compounds of the present invention to a protein, the R groups may be reacted with a protein to produce a protein conjugate. Preferred proteins are those in serum, wherein the compound of the invention is directly injected and the conjugate is formed *in situ*. It is understood that other functional groups, as described above, may be used to link the bifunctional metal-chelating ligands of this invention to proteins such as monoclonal antibodies or fragments thereof.

- 19 -


Compounds of the formula I can generally be prepared as follows:

Ligands in which the aza macrocyclic ring carbon atoms are modified are built de novo from  
5 suitable aziridine precursors by cyclotetramerization.

Cyclotetramerization of N-benzylaziridine has been reported in the literature. (See, for example, T. E. Burg and G. R. Hansen, J. Heterocyclic  
10 Chemistry, (1968), 5, 305.) The synthetic approach to the preparation of S,S,S,S-tetramethyltetra-azacyclododecane (5) is given in scheme 1:

- 20 -

SCHEME 1



- 21 -

[S]-N-Benzoylalanine(1) is reduced with diborane to yield [S]-N-benzylalaninol(2). Under Mitsunobu conditions, compound (2) affords [S]-N-benzyl-2-methyl aziridine(3). Cyclotetramerization under p-toluene sulfonic acid catalysis in ethanol, followed by treatment with ammonium hydroxide (NH<sub>4</sub>OH), furnishes S,S,S,S-tetra-N-benzyl-tetramethyltetraazacyclododecane(4), which is debenzylated under transfer hydrogenolytic conditions to obtain S,S,S,S-tetramethyltetraazacyclododecane(5).

Tetraalkylation of compound (5) with t-butyl bromoacetate in the presence of sodium carbonate, followed by deprotection with trifluoroacetic acid and anisole, affords the ligand (6).

Tetraalkylation of compound (5) with benzyl 2-triflyloxylactate (prepared as described by S.I. Kang et al., Inorg. Chem., (1993), 32, 2912-2918) in the presence of sodium carbonate, followed by catalytic hydrogenolytic debenzylation furnishes the ligand (7).


Tris-alkylation of compound (5) with t-butyl bromoacetate in the presence of sodium bicarbonate, followed by deprotection by treatment with trifluoroacetic acid and anisole, provides the ligand (8).

For preparing multimeric ligands, 1,4,7-tris-carboxymethyl-1,4,7,10-tetraazacyclododecane-10-yl (DO3A) units, for example, are attached to a bridging unit by two different methods. In the first method, a spiro epoxide on the bridging unit is used to alkylate DO3A. In the second method, a glycidyloxy moiety is attached to the bridging unit which then is used to alkylate DO3A.

- 22 -


By way of example, N-alkylation of DO3A with suitable epoxides of formula (9) below, wherein X is a carbocyclic, fused heterocyclic or spiro-heterocyclic unit, and n is 2, 4 or 8, affords

5 multimeric ligands of the formula (10):




Epoxides of formula (9) are generated by the

10 Prilezhaev Reaction in which olefins of formula (11) are treated with peracids such as m-chloroperbenzoic acid:



Olefins of formula (11) are generated by the

alkylation of alcohols of formula (12):




Another mode of attachment of DO3A would be to epoxidize olefins of formula (13) to obtain epoxides




- 23 -


of formula (14), and then alkylating DO3A with the epoxides to form ligands of formula (15):




These two method of attachment of DO3A could also be present in the same bridging unit X leading to structures as in formula (16):




Representative bridging units are exemplified in formulas (17), (18), (19), (20), (21) and (22):




- 24 -



20




21



22


The structures shown in formulas (17) to (22) are only by way of representative examples. They will, in actuality, be a mixture of all the possible diastereomers, if the attachment is through a 2-hydroxypropyloxy link. In the case of formula 22, for example, in addition to the presence of diastereomers, the product would also consist of the various geometric isomers that could result when the two myo-inositol molecules are coupled to each other. The coupling of the two myo-inositol units is achieved by reacting the compound of formula (23) with cyclohexane-1,4-dione (24):

- 25 -



The bridging unit for formula 17 is generated from commercially available diethyl cyclohexane-1,4-dione-2,5-dicarboxylate by lithium aluminum hydride reduction as described in J.G. Murphy, J. Med. Chem., (1966), 9, 157.

The bridging unit for formula (18) is generated by treating myo-inositol (25) with 2,2-dimethoxypropane in the presence of p-toluene sulfonic acid as described by Giggs et al., Carbohydrate Res., 1985, 142, 132:



25

15

The bridging unit (19) is made from (18) by acidic hydrolysis. The unit (20) is prepared from myo-inositol (25) by reaction with excess 2,2-dimethoxypropane in the presence of p-toluene-sulfonic acid. The unit (21) is made from the bis-


- 26 -

epoxide precursor to the unit (17) by methods described above for functionalizing hydroxy groups.

The ligand of Example (10), below, bearing a phosphonomethyl arm, is made by treating DO3A with phosphorus acid and formaldehyde as described by M. Tazaki et al., Chem. Lett., 1982, 571.

All stereoisomers of the compounds and complexes of the present invention are contemplated herein, whether alone (that is substantially free of other isomers), in a mixture of certain stereoisomers (for example, as a racemate) or in any other mixture thereof.


The invention will now be further described by the following examples. In the structures of examples 1 - 6, R is



- 27 -

Example 1

(1 $\alpha$ , 2 $\alpha$ , 4 $\beta$ , 5 $\beta$ )-10,10'-[(1,2,4,5-Tetrahydroxy-1,4-cyclohexanediyl)bis(methylene)]bis[1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid],  
digadolinium salt



10 A. trans-2,5-dihydroxy-1,4-dimethylene cyclohexane

A warm solution of dry, recrystallized diethyl 1,4-cyclohexanedione-2,5-dicarboxylate (30.75 g, 120 mmol) in 350 ml anhydrous tetrahydrofuran (THF) was added to a refluxing solution of 1.0 M LAH in THF (600 mL) over a 2 hour period. After an additional 1.5 hours at reflux and subsequent cooling, a saturated solution of aqueous Rochelles's salt (app. 73 mL) was added dropwise to the reaction mixture. The tartrate complex was filtered, washed with THF and the filtrate was concentrated to a paste. The paste was recrystallized from hot ethyl acetate; yield 4.39 g (31.3 mmol, 26.1%). A second recrystallization yielded 3.81 g (27.2 mmol, 22.6%). M.P.: 160 - 161.5°C, uncorrected (lit. 162.5 - 163°C).

B. (3 $\alpha$ , 4 $\alpha$ , 6 $\beta$ , 9 $\beta$ )-1,7-dioxadispiro[2.2.2.2]-decane-4,9-diol

A solution of wet, 80% *m*-chloroperoxybenzoic acid (mCPBA, 7.00 g, app. 30 mmol) in 30 mL dry dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>) was dried over anhydrous

- 28 -

magnesium sulfate ( $\text{MgSO}_4$ ). This solution was added to a solution of dry Compound A (1.40 g, 10.0 mmol) in 15 mL dry  $\text{CH}_2\text{Cl}_2$ ). The mixture was stirred at room temperature under a dry nitrogen atmosphere for 16 hours. A second sample (550 mg) of mCPBA was added to ensure complete conversion of the dimethylene compound to the diepoxide. After 3 hours, the mixture was evaporated to dryness and excess mCPBA/*m*-chlorobenzoic acid was removed by treating the residue with diethyl ether. The ether-insoluble material was filtered, washed with ethyl ether ( $\text{Et}_2\text{O}$ ) and air dried to yield 1.35 g of crude Compound B. The crude material was recrystallized from a minimum of hot methanol; yield 967 mg (5.6 mmol, 56.1%).  
M.P.: 231 - 233°C.

C.  $(1\alpha, 2\alpha, 4\beta, 5\beta)$ -10,10'-[(1,2,4,5-Tetrahydroxy-1,4-cyclohexanediyl)bis(methylene)]bis[1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid], hexasodium salt

The pH of a suspension of 1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid (DO3A)- $\text{H}_2\text{SO}_4$  (prepared as described in D.D. Dischino et al., Inorg. Chem., 1991, 30, 1265 - 69; 2.67 g, 6.0 mmol) in 2.0 mL water was adjusted to approximately pH 12.5 with 5.40 mL 5N sodium hydroxide. This solution was warmed to 50°C, and Compound B (517 mg, 3.0 mmol) was added portion-wise over 5.5 hours. Crystals developed overnight from the warm mixture. After cooling, the crude crystals (3.4 g) were isolated and were recrystallized from 20 mL hot water to yield 1.3 g (1.4 mmol, 45%) of Compound C.

- 29 -

Mass Spectrum (FAB): (M-7H+8Na)<sup>+</sup> at 1040; (M-6H+7Na)<sup>+</sup> at 1019; (M-5H+6Na)<sup>+</sup> at 997; (M-4H+5Na)<sup>+</sup> at 975; (M-3H+4Na)<sup>+</sup> at 953; (M-2H+3Na)<sup>+</sup> at 931.

Elemental analysis: Found: C 42.53, H 6.10, N

5 10.90. Calculated for: C<sub>36</sub>H<sub>58</sub>N<sub>8</sub>Na<sub>6</sub>O<sub>16</sub>·1.16 H<sub>2</sub>O: C 42.49, H 5.97, N 11.01.

D. (1 $\alpha$ ,2 $\alpha$ ,4 $\beta$ ,5 $\beta$ )-10,10'-[1,2,4,5-Tetrahydroxy  
-1,4-cyclcohexanediyl)bis(methylene)]  
10 bis[1,4,7,10-tetraazacyclododecane-1,4,7-  
triacetic acid].digadolinium salt

The pH of a solution of Compound C (760 mg, 762  $\mu$ mol) in 5.0 mL deionized water was adjusted to pH 4.83 with 225  $\mu$ L acetic acid (HOAc). A solution  
15 of Gd(OAc)<sub>3</sub>·4 H<sub>2</sub>O (680 mg, 1.67 mmol) in 5.0 mL warm deionized water was added to the ligand solution over a 10 minute period; final pH 4.73. After 1 minute of stirring at room temperature, a precipitate began to develop. The volume of the precipitate increased  
20 with time and the reaction was left to run overnight. The precipitate was filtered, extensively washed with water, and dried; yield 741 mg (631  $\mu$ mol, 83%). An analytical sample was prepared by recrystallization from hot water.

25 Mass spectrum: (FAB): (M+H)<sup>+</sup> at 1169 through 1177.

Elemental analysis:

Found: C 35.93, H 5.50, N 8.84.

Calculated for C<sub>36</sub>H<sub>58</sub>Gd<sub>2</sub>N<sub>8</sub>O<sub>16</sub>·2.5 H<sub>2</sub>O: C 35.49,


H 5.21, N 9.20.

30

- 30 -

Example 2

(3 $\alpha$ , 4 $\alpha$ , 5 $\beta$ , 6 $\alpha$ , 7 $\beta$ , 7 $\alpha\alpha$ )-10,10'-10'',10'''-[[Hexahydro-2,2-dimethyl-1,3-benzodioxol-4,5,6,7-tetrayl]tetra(oxy)-  
 5 tetra(2-hydroxy-3,1-propanediyl)]tetrakis[1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid],  
tetragadolinium salt



10

A. (+)-1,2-O-Isopropylidene-myo-inositol

A mixture of myo-inositol (25.0 g, 140 mmol),  
 2,2-dimethoxypropane (42.5 mL, 350 mmol) and p-  
 toluenesulfonic acid (250 mg) in dimethylsulfoxide  
 15 (80 mL) was stirred at 100°C for 45 minutes until the  
 solution became homogenous. After stirring the  
 mixture at ambient temperature for 30 minutes,  
 triethylamine (2.5 mL), ethanol (100 mL) and ether  
 (500 mL) were introduced causing precipitation. The  
 20 solid material was filtered, washed with ether-  
 methanol (200 mL, 5:1) and ether, and dried to obtain  
 the ketal compound A (9.72 g) as a white solid in 32%  
 yield.

m.p.: 182.4 - 182.9°C

25



- 31 -

B. (3 $\alpha$ ,4 $\alpha$ ,5 $\beta$ ,6 $\alpha$ ,7 $\alpha$ ,)-Hexahydro-2,2-  
dimethyl-tetra-kis-4,5,6,7-(2-propenyl-  
oxy)-1,3-benzodioxole

To a suspension of sodium hydride (545 mmol)  
5 in dimethylformamide (50 mL) was added the ketal  
compound A (20.0 g, 90.8 mmol) dimethylformamide (20  
mL) at ambient temperature under nitrogen. After  
stirring the suspension for 2 hours, allyl bromide  
(83.4 g, 689 mmol) in dimethylformamide (50 mL) was  
10 reacted for 18 hours. The unreacted sodium hydride  
was carefully quenched with water, and then the  
solvent was removed to obtain a dark brown oil. The  
residue was partitioned in water (100 mL) and  
dichloromethane (100 mL) and extracted with  
15 dichloromethane (100 mL x 3). The combined organic  
layers were dried over sodium bisulfate and  
evaporated to obtain a brownish liquid (about 45 mL).  
The crude product was purified by silica gel flash  
chromatography (ethyl acetate and hexanes, 1:8) to  
20 afford compound B (26.2 g) as a pale yellow liquid in  
76% yield.

C. (3 $\alpha$ ,4 $\alpha$ ,5 $\beta$ ,6 $\alpha$ ,7 $\alpha$ ,)-Hexahydro-2,2-  
dimethyl-tetra-kis-4,5,6,7-(oxiranylmethoxy)-  
25 1,3-benzodioxole

To compound B (6.0 g, 15.8 mmol) in  
dichloromethane (10 mL) was added dropwise m-  
chloroperbenzoic acid (85% from ICN, 19.2 g, 94.6  
mmol) dissolved in dichloromethane (100 mL). A  
30 homogeneous solution was initially obtained but a  
white solid, presumed to be m-chlorobenzoic acid,  
slowly precipitated out of the solution after about  
15 minutes. After 24 hours, sodium metabisulfite  
(100 mL, 5% in water) and sodium bicarbonate (200 mL,

- 32 -

10% in water) were added and the resulting solution was extracted with dichloromethane (100 mL x 3). The organic layers were combined and dried over sodium sulfate. The residue, obtained from removal of the solvent, was purified by flash silica gel chromatography to obtain compound C as a colorless liquid in 76% yield.

D. (3 $\alpha$ ,4 $\alpha$ ,5 $\beta$ ,6 $\alpha$ ,7 $\beta$ ,7 $\alpha$ ,)-  
10 10,10',10'',10'''-[[Hexahydro-2,2-dimethyl-  
1,3-benzodioxol-4,5,6,7-tetrayl]-  
tetra(oxy)tetra(2-hydroxy-3,1-  
propanediyl)]tetrakis[1,4,7,10-tetraazacyclo-  
dodecane-1,4,7-triacetic acid], triethylamine  
15 (1:4) salt

To a suspension of DO3A (9.77 g, 28.2 mmol) dissolved in alkaline water (6 mL, 10 N sodium hydroxide was used to obtain a pH 10 solution) at 40-50°C was added the tetraepoxide compound C (1.54 g, 20 3.46 mmol) in acetonitrile (5 mL). The reaction mixture was stirred for 2 days maintaining the same pH. The crude product was purified by DEAE Sephadex chromatography using triethylammonium bicarbonate (pH 7.5) as the eluent. The buffer concentration to 25 bring out the product compound D was 200 to 500 mM. Removal of the buffer gave compound D (7.8 g) in 58% yield.  
m.p.: 178°C (decomp.)

- 33 -


E. (3 $\alpha$ , 4 $\alpha$ , 5 $\beta$ , 6 $\alpha$ , 7 $\beta$ , 7 $\alpha\alpha$ )-10,10'-10'',10'''-  
[[Hexahydro-2,2-dimethyl-1,3-benzodioxol-  
4,5,6,7-tetrayl]tetra(oxy)tetra(2-hydroxy-  
3,1-propanediyl)]tetrakis[1,4,7,10-  
5 tetraazacyclododecane-1,4,7-triacetic acid],  
tetragadolinium salt

To compound D (216 mg, 96  $\mu$ mol) dissolved in  
water (1 mL) was added gadolinium acetate (236 mg,  
580  $\mu$ mol) dissolved in water (2 mL). The solution  
10 was heated to 65°C for 18 hours. The pH was  
maintained between 4.0 and 6.0 during the chelation  
reaction. The material was loaded to a CHP20P column  
and eluted with water (1.9 L) followed by 10%  
ethanolic water (1 L). The product was eluted with  
15 10% ethanol-containing fraction, as identified by  
HPLC, and evaporated to dryness to yield the title  
compound (283 mg) as a white crystalline solid in a  
quantitative yield.  
MS (FAB): m/z: 2447.8 [(M+H)<sup>+</sup>, base peak]. Analysis  
20 Calculated for C<sub>77</sub>H<sub>124</sub>N<sub>16</sub>O<sub>34</sub>Gd<sub>4</sub>•11.41H<sub>2</sub>O: C, 34.87;  
H, 5.58; N, 8.45.  
Found: C, 34.68; H, 5.98; N, 8.30; H<sub>2</sub>O, 7.75  
(desorption KF).

- 34 -

Example 3

3,4,5,6-Tetra-O-[2-hydroxy-3-[4,7,10-  
 tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-  
 5 yl]propyl]-myo-inositol, tetragadolinium salt



A. 3,4,5,6-Tetra-O-[2-hydroxy-3-[4,7,10-tris-  
 10 (carboxymethyl)-1,4,7,10-tetraazacyclo-  
 dodecan-1-yl]propyl]-myo-inositol,  
tetratriethyl ammonium salt

Compound D from Example 2 was treated with 1.0  
 N aqueous hydrochloric acid (10 mL) for 0.5 hours to  
 15 remove the ketal group. The resulting solution was  
 applied to a polyvinylpyridine (PVP) column (2.5 x 30  
 cm) and eluted with water. Silver nitrate was given  
 to test each fraction to detect any breakage of  
 chloride ion. Removal of the water from the  
 20 fractions containing the product gave the title  
 compound (910 mg) in 80% yield.

m.p.: 210°C (decomp). MS (FAB): m/z: 1789.9  
 [(M+H)<sup>+</sup>, base peak]; 1811.9 [(M+Na)<sup>+</sup>]. Analysis  
 Calculated for C<sub>74</sub>H<sub>132</sub>N<sub>16</sub>O<sub>34</sub>·11.0H<sub>2</sub>O:

25 C, 44.63; H, 7.81; N, 11.27. Found: C, 44.63; H,  
 7.84; N, 11.41; H<sub>2</sub>O, 6.43 (desorption KF); ROI, 0.17.

- 35 -

B. 3,4,5,6-Tetra-O-[2-hydroxy-3-[4,7,10-  
tris(carboxymethyl)-1,4,7,10-tetraazacyclo-  
dodecan-1-yl]propyl]-myo-inositol,  
tetragadolinium salt


5 To gadolinium acetate (595 mg, 1.46 mmol)  
dissolved in water (1.5 mL) was added Compound A (550  
mg, 0.246 mmol) dissolved in water (2 mL). The  
chelation mixture was kept overnight at 60°C and the  
pH of the solution maintained between 4.0 and 5.0.  
10 The resulting mixture was loaded to a CHP20P column  
(2.5 x 25 cm) and eluted with water followed by 10%  
ethanol. The title compound was brought out by 10%  
ethanolic water. Removal of the solvents from the  
fractions containing the product gave the title  
15 compound as a white solid in 93% yield.  
MS (FAB): 2408.7[(M+H)<sup>+</sup>, <sup>158</sup>Gd]): Analysis Calculated  
for C<sub>74</sub>H<sub>120</sub>N<sub>16</sub>O<sub>34</sub>Gd<sub>4</sub>•4.86 H<sub>2</sub>O: C, 35.63; H, 5.24; N,  
8.98. Found: C, 35.48; H, 5.74; N, 8.75; 3.51 H<sub>2</sub>O  
(desorption KF); ROI, 31.52.

20

- 36 -

Example 4

(1 $\alpha$ , 2 $\alpha$ , 3 $\alpha$ , 4 $\beta$ , 5 $\alpha$ , 6 $\beta$ )-10,10'-[(2,3,5,6-tetrahydroxy-1,4-cyclohexanediyl)bis(2-hydroxy-3,1-propanediyl)]-bis[1,4,7-10-tetraazacyclododecane-1,4,7-triacetic acid], digadolinium salt



10 A. [3aR-(3a $\alpha$ , 4 $\alpha$ , 4a $\alpha$ , 7a $\alpha$ , 8 $\beta$ , 8b)]-Hexahydro-2,2,6,6-tetramethyl-4,8-bis-(2-propenyl-oxv)benzo[1,2-d:4,5-d']bis[1,3]dioxole

A sample of sodium hydride (NaH) (8.31 g, 156 mmol; 45% in mineral oil) was added to an ice cold solution of dry 1,2:4,5-di-O-isopropylidene-myoinositol (10.00 g, 38.4 mmol, prepared as indicated in Gigg et al., Carbohydr. Res., 1985, 142, 132) in dry dimethylformamide (DMF) (80 ml). After stirring for 0.5 hours at room temperature, allyl bromide (8.31 mL, 96 mmol) was added dropwise over 5 minutes. During the addition, the temperature was maintained at 25°C by cooling in a cold water bath. After 1.5 hours, the reaction was stopped by the careful addition of water and the mixture was evaporated to dryness in vacuo. The residue was dissolved in water, extracted with ethyl acetate (EtOAc) (4 x 50 ml), and washed with water (2 x 20 ml) and brine (1 x 20 ml). The organic layer was dried over sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to dryness under reduced pressure. The residue was

- 37 -

recrystallized from hexane to give pure product as a colorless solid (12.88 g, yield 98.5%).

M.P. 82 - 83°C, uncorrected.

- 5 B. (3 $\alpha$ ,4 $\alpha$ ,4 $\alpha$ ,7 $\alpha$ ,8 $\beta$ ,8 $\beta$ )-Hexahydro-2,2,6,6-tetramethyl-4,8-bis(oxiranylmethoxy)benzo  
[1,2-d:4,5-d']bis[1,3]dioxole

A solution of 80% mCPBA (13.09 g, 60.7 mmol) in dry dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>) (95 ml) was added to  
10 a cooled solution containing Compound A (5.11 g, 15.0 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (15 ml). After stirring at room temperature overnight, the mixture was treated with saturated aqueous sodium sulfite (Na<sub>2</sub>SO<sub>3</sub>) followed by saturated aqueous sodium bicarbonate (NaHCO<sub>3</sub>) until  
15 the aqueous layer was neutral. The organic layer was dried over sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>) followed by evaporation to dryness in vacuo. The residue was recrystallized from hot ethanol to give the pure product (4.05 g, yield 72.4%).  
20 M.P. 103 - 105°C (uncorrected)

- C. (1 $\alpha$ ,2 $\alpha$ ,3 $\alpha$ ,4 $\beta$ ,5 $\alpha$ ,6 $\beta$ )-10,10'-[(2,3,5,6-tetrahydroxy-1,4-cyclohexanediyl)bis(2-hydroxy-3,1-propanediyl)]bis[1,4,7-10-tetraazacyclododecane-1,4,7-triacetic acid], triethylamine  
25 (1:2) salt

A sample of crude (3 $\alpha$ ,5 $\alpha$ ,5 $\beta$ ,6 $\alpha$ ,7 $\beta$ ,7 $\alpha$ )-10-10'-[[hexahydro-2,2-dimethylbenzol[1,2-d:4,5-d']-bis[1,3]dioxol-4,8-diyl]di(oxy)di(2-hydroxy-3,1-propanediyl)]bis[1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid] was prepared by heating a  
30 solution of DO3A•H<sub>2</sub>SO<sub>4</sub> (28.24 g, 63.5 mmol) and Compound B (5.92 g, 15.9 mmol) in 10 M sodium hydroxide (NaOH) (28.0 mL), H<sub>2</sub>O (20 mL) and

- 38 -

1,4-dioxane (17 mL) at 50°C for 70 hours. The pH of the crude mixture was adjusted to pH 1 using 1N hydrochloric acid (HCl) and was heated at 55°C for 1.5 hours. After the hydrolysis was complete, the pH  
5 of the solution was adjusted to 9 with dilute NaOH and the mixture was applied to a DEAE-Sephadex A-25 column ( $\text{HCO}_3^-$  form, 2L). The column was washed with water, and the product was eluted with a linear gradient of 5 - 250 mM TEAB buffer (pH 7.5, 4 L  
10 each). The fractions which contained pure Compound C were combined, freed of TEAB, and lyophilized to give Compound C as the di(triethylammonium) salt (18.22 g, yield 97.2%). The mass spectral analysis of this product indicated the presence of chloride ion. A  
15 sample of the contaminated product (10.00 g) was dissolved in water and was applied onto a AG50W x 8 column ( $\text{H}^+$  form, 500 ml). The column was washed with water, and the product was eluted with 0.5 M ammonium hydroxide ( $\text{NH}_4\text{OH}$ ). The fractions which contained  
20 Compound C were combined and evaporated to give the ammonium salt (2.91 g). The ammonium salt was dissolved in water, applied to a DEAE-Sephadex A-25 column ( $\text{HCO}_3^-$  form, 2L). The column was washed with water and the product was eluted with 200 mM TEAB (pH  
25 7.5, 5L). The fractions containing pure material were combined, freed of TEAB, and lyophilized to give pure Compound C as the di(triethylammonium) salt (2.86 g, theo. yeild 25.9%).



- 39 -

- D. (1 $\alpha$ ,2 $\alpha$ ,3 $\alpha$ ,4 $\beta$ ,5 $\alpha$ ,6 $\beta$ )-10,10'-[(2,3,5,6-tetrahydroxy-1,4-cyclohexanediyl)]bis(2-hydroxy-3,1-propanediyl)]bis[1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid], digadolinium salt

A sample of Compound C (1.02 g) was dissolved in a solution containing Gd(OAc)<sub>3</sub>•4 H<sub>2</sub>O (1.219 g, 3 mmol) in water (10 mL) and was heated at 50°C for 16 hours. The solution was applied onto a CHP20-P column (500 mL). The salt contaminants were eluted with water, and the product was eluted with 10% ethanol in water. The appropriate fractions were combined and evaporated to dryness in vacuo. The glassy residue was dissolved in water, and lyophilized to give pure title compound (1.04 g, yield 80.5%).

MS (FAB): (M + H)<sup>+</sup> at 1295.2 (major isotope).

Elemental analysis:

Found: C, 36.83; H, 5.63; N, 8.46%.

20 Calculated for C<sub>40</sub>H<sub>66</sub>Gd<sub>2</sub>N<sub>8</sub>O<sub>20</sub>•1.20 H<sub>2</sub>O:


C, 36.53; H, 5.24; N, 8.52%.

- 40 -

Example 5

10-[[1,4-Dihydroxy-2,5-bis[2-hydroxy-3-[4,7,10-tris-  
(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-yl]-  
5 propoxy]-4-[[4,7,10-tris(carboxymethyl)-1,4,7,10-  
tetraazacyclododecan-1-yl]methyl]cyclohexyl]methyl]-  
1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid,  
tetragadolinium salt

10



A. 1,7-dioxadispiro [2,2,2,2]-decane-4,9-diallyldiol

To a suspension of sodium hydride (432 mg, 18  
15 mmol) in anhydrous dimethylformamide (15 ml), a  
solution of bisepoxydiol (1.0 g, 6 mmol) (described  
in Example 1B) in anhydrous dimethylformamide (60 ml)  
was added dropwise and the mixture was stirred at  
room temperature for 60 minutes. Allyl bromide  
20 (2.178 g, 18 mmol) was added and the mixture was  
stirred at room temperature for 18 hours. Excess of  
sodium hydride was decomposed with water (2 ml),  
dimethylformamide removed in vacuo and the residue  
extracted with ethyl acetate (2 x 100 ml), washed  
25 with water (2 x 50 ml) and dried. Solvent removal  
afforded the crude product. Silica gel (25 g) column  
chromatography using hexanes and ethyl acetate (2/1)  
as the eluent afforded the pure diallyl derivative  
compound A as a colorless solid (0.7 g, yield 48%).  
30 Melting point: 63.5-65.5°C (uncorrected).

- 41 -

B. (3 $\alpha$ ,4 $\alpha$ ,6 $\beta$ ,9 $\beta$ )-4,9-bis(oxiranylmethoxy)-  
1,7-dioxadispiro[2.2.2.2]decane

A solution of m-chloroperoxybenzoic acid (4.3 g, 80%, 18.5 mmol) in dichloromethane (30 ml) was  
5 dried over anhydrous magnesium sulfate. This solution was added dropwise to a solution of compound A (1.25 g, 5 mmol) in dry dichloromethane (25 ml). The reaction mixture was stirred at room temperature for 20 hours. Excess of m-chloroperoxybenzoic acid  
10 was decomposed with saturated aqueous sodium metabisulfate solution (15 ml). The organic layer was separated, washed with saturated sodium bicarbonate (2 x 50 ml), and with water (2 x 50 ml). The organic layer was dried over magnesium sulfate  
15 and evaporated to dryness. Recrystallization of the crude material from hot ether afforded the pure tetraepoxide compound B as white crystals (1.1 g, yield 77.5%).  
Melting point: 93.0-96.0°C (uncorrected).

20

C. 10-[[[1,4-Dihydroxy-2,5-bis[2-hydroxy-3-  
[4,7,10-tris-(carboxymethyl)-1,4,7,10-  
tetraazacyclododecan-1-yl]-propoxy]-4-  
25 [[4,7,10-tris(carboxymethyl)-1,4,7,10-  
tetraazacyclododecan-1-  
yl)methyl]cyclohexyl)methyl]-1,4,7,10-  
tetraazacyclododecane-1,4,7-triacetic acid,  
triethylamine (1:4) salt

A solution of DO3A (17.3 g, 50 mmol) in water  
30 was made (80 ml) and the pH of the solution adjusted to 12 with 5 N sodium hydroxide solution. The solution was heated to 80°C and a solution of compound B (1.42 g, 5 mmol) in dioxane (10 ml) was added dropwise. The mixture was stirred at this

- 42 -

temperature for 82 hours. At the end of 82 hours, the pH of the reaction mixture was adjusted to 7 with acetic acid. The mixture was diluted and loaded on to a Sephadex G-25 column. The column was eluted first with 100-250 mM triethyl ammonium bicarbonate (TEAB) buffer and then with 250-500 mM TEAB buffer. Fractions containing the pure tetrameric material were combined and solvent removal afforded the pure product as the tetra triethyl ammonium salt of the title compound as a colorless glassy solid (2.85 g, yield 35%).

Mass Spectrum: 1670 (M+H)<sup>+</sup>. 102 (CH<sub>3</sub>CH<sub>2</sub>)<sub>3</sub>N<sup>+</sup>H

Elemental Analysis: Found: C, 53.70; H, 9.15; N, 13.62; H<sub>2</sub>O, 0.58%. Calculated for C<sub>94</sub>H<sub>184</sub>N<sub>20</sub>O<sub>30</sub>·0.67 H<sub>2</sub>O: C, 54.11; H, 8.85; N, 13.42; O, 23.52%.

D. 10-[[[1,4-Dihydroxy-2,5-bis[2-hydroxy-3-[4,7,10-tris-(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-yl]-propoxy]-4-[[4,7,10-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-yl]methyl]cyclohexyl]methyl]-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, tetragadolinium salt

25 A sample of Compound C (0.2 g) was dissolved in 3 mL deionized water (reactant I). In a separate vial, 0.18 g of GdCl<sub>3</sub> powder was dissolved in 5 mL deionized water (reactant II). At about 70°C and constant stirring, reactant I was added into reactant II dropwise over a time period of several hours. Meanwhile, 5N NaOH was pipetted into the solution from time to time to neutralize the protons released from chelation and maintain the reaction pH around 6. The excess metal ions were precipitated in the form

- 43 -

of  $M(OH)_3 \cdot x H_2O$  by raising solution pH to about 9 and incubating at both about 70°C and room temperature for several hours. The precipitate was subsequently removed through centrifugation and  
5 filtration with 0.22  $\mu m$  membrane. The filtrate was finally condensed and neutralized to pH 7 in preparation for HPLC purification.

The crude chelate product was purified by preparative HPLC, using mobile phase gradient  
10 (acetonitrile:water) and silica-based reverse phase column (YMC C18, 5 $\mu$ , 200 Å). The solvent was removed from the fractions containing the product. The residue was further dried in a vacuum oven at about 70°C overnight. The yield of the desired chelate was  
15 70%.

Mass Spectrum (FAB, m/e):  $(Gd^{159} + H)^+$  at 2287.5.


Elemental Analysis (C,H,N):

Calculated for  $C_{70}H_{112}N_{16}O_{30}Gd_4 \cdot 7.73 H_2O$ : C 34.66,  
H 5.30, N 9.24%.

20 Found: C 34.26, H 5.36, N 9.08%.

Example 6

[3aR-(3a $\alpha$ , 3"a $\alpha$ , 4 $\beta$ , 4" $\beta$ , 5 $\alpha$ , 5"a $\alpha$ , 6 $\beta$ , 6" $\beta$ , 7 $\alpha$ , -  
 7" $\alpha$ , 7a $\alpha$ , 7"a $\alpha$ )]-dodecahydro-4,4",5,5",6,6",7,7"-  
 5 octakis-[[2-hydroxy-3-[(4,7,10-tricarboxymethyl)-  
 1,4,7,10-tetraazacyclododecan-1-yl]propyl]-  
 oxy]dispiro-[1,3-benzodioxole-2,1'-cyclo-hexane-  
 4',2"-[1,3]-benzodioxole], octagadolium salt

A. (+)-3,4,5,6-Tetra-O-allyl-myo-inositol

Compound B from Example 2 (15.3 g, 44.9 mmol)  
 was dissolved in methanol (30 mL) and treated with 3N  
 15 hydrochloric acid (30 mL), with constant stirring at  
 room temperature for 13 hours. The reaction mixture  
 was neutralized with saturated sodium bicarbonate  
 solution and extracted with methylene chloride (50 mL  
 x 3). The dichloromethane layers were combined and  
 dried with sodium sulfate, filtered, evaporated in  
 20 vacuo and purified by column chromatography (silicon  
 dioxide, ethyl acetate: hexanes (1:1 v/v)) to obtain  
 9.1 g (67%) of the desired product compound A.

- 45 -

B. [3aR-(3a $\alpha$ ,3\*a $\alpha$ ,4 $\alpha$ ,4\* $\beta$ ,5 $\beta$ ,5\*a $\alpha$ ,6 $\alpha$ ,6\* $\beta$ ,  
7 $\beta$ ,7\*a $\alpha$ ,7a $\alpha$ ,7\*a $\alpha$ )]-Dodecahydro-4,4\*,5,-  
5\*,6,6\*,7,7\*-octakis-(2-propenyloxy)-  
dispiro[1,3-benzo-dioxole-2,1'-cyclo-  
5 hexane-4',2'-[1,3]-benzodioxole]

A mixture of the diol compound A (1.0 g, 2.94 mmol), 1,4-cyclohexanedione (165 mg, 1.47 mmol) and p-toluenesulfonic acid (55 mg, 0.289 mmol) was heated in toluene (30 mL) at reflux for 19 hours. The crude  
10 reaction mixture was purified by flash silica gel column chromatography (about 200 g) using different eluents of hexane and ethyl acetate (2:1, 1:1 and 1:2, 500 mL each). Three to four spots that have the molecular ion peak at 757 (m/z) were pooled to afford  
15 an oil (520 mG) in 43% yield, and the resulting mixture used as a mixture of the isomers of the expected octa-allyl bis-ketal product.  
TLC: R<sub>f</sub> 0.45, 0.37 and 0.32 in hexanes and : acetone (5:1, v/v)

20

C. [3aR-(3a $\alpha$ ,3\*a $\alpha$ ,4 $\beta$ ,4\* $\beta$ ,5 $\alpha$ ,5\*a $\alpha$ ,6 $\beta$ ,6\*b,  
7 $\alpha$ ,7\*a $\alpha$ ,7a $\alpha$ ,7\*a $\alpha$ )]-Dodecahydro-  
4,4\*,5,5\*,6,6\*,7,7\*-octakis-(oxyranylmethoxy)-  
dispiro[1,3-benzodioxole-2,1'-cyclohexane-  
25 4',2'-[1,3]-benzodioxole]

The oxidation of the octa-allyl compound B (1.13 g, 1.49 mmole) by m-chloroperoxybenzoic acid (3.0 g, 85%, 17.6 mmole) in dichloromethane (20 mL) was carried out at ambient temperature for 50 hours.  
30 The white solid, which precipitated out, during the oxidation was identified as m-chlorobenzoic acid by <sup>1</sup>H-NMR (CDCl<sub>3</sub>). After removal of the solid by filtration, the dichloromethane filtrate was treated with sodium metabisulfite and sodium hydroxide where

- 46 -

the final pH of the aqueous layer was 11.7. The organic layer was dried over sodium sulfate and concentrated in vacuo. The crude solid product was purified by column chromatography (flash silica gel, 75 g) to obtain a white solid (670 mg) in 51% yield.

D. [3aR-(3a $\alpha$ ,3\*a $\alpha$ ,4 $\beta$ ,4\* $\beta$ ,5 $\alpha$ ,5\*a $\alpha$ ,6 $\beta$ ,6\*b,7 $\alpha$ ,7\*a,7a $\alpha$ ,7\*a $\alpha$ )]-dodecahydro-4,4",5,5",-6,6",7,7"-octakis-[[2-hydroxy-3-[(4,7,10-tricarboxymethyl)-1,4,7,10-tetraazacyclo-dodecan-1-yl]propyl]oxy]dispiro-[1,3-benzodioxole-2,1'-cyclohexane-4',2'-[1,3]-benzodioxole]

The octaepoxide Compound C (200 mg, 235  $\mu$ mol) in acetonitrile (0.5 mL) was added to DO3A (2.41 g, 7.04 mmol) in water (4-5 mL) whose pH was adjusted to 9.6 by 10 N sodium hydroxide at 60°C. The resulting solution was kept for 48 hours and analyzed by PRP-X 100 HPLC column at desirable intervals. Two relatively pure fractions were obtained after two DEAE Sephadex A-25 ion exchange columns. One eluted by 300-400 mM triethyl-ammonium bicarbonate (TEAB) showed a peak at m/z 3328.0 corresponding to the heptamer, whereas the other by 400 mM TEAB turned out to be the expected octamer with a peak at m/z 3656.6 (C<sub>154</sub>H<sub>268</sub>N<sub>32</sub>O<sub>68</sub>).

<sup>1</sup>H-NMR (D<sub>2</sub>O):  $\delta$  1.15 (t, 106H, CH<sub>3</sub>CH<sub>2</sub>N); 1.7-1.9 (m, methylenes of the middle cyclohexane, 8H); 3.07 (q, 71H, CH<sub>3</sub>CH<sub>2</sub>N), 2.8-4.6 (m, 228H, all of methines and methylenes of the octamer ligand except the triethylamine and bridged cyclohexane). <sup>13</sup>C-NMR (D<sub>2</sub>O): 8.10 and 46.29 (CH<sub>3</sub>CH<sub>2</sub>N of triethyl-amine); 41.88, 46.29, 49.25, 49.54, 49.62, 49.76, 50.04,



- 47 -

55.72, 56.43, 58.69, 172 - 175 (broad due to 24 carboxylate and/or carboxylic acid groups).

MS (FAB): 3,328 (M+H)+.

Analysis Calculated for  $C_{154}H_{268}N_{32}O_{68}$ -

5  $8[N(C_2H_5)_3] \cdot 17.20H_2O$ :

C, 50.81; H, 8.92; N, 11.73. Found: C, 50.33; H, 8.92; N, 11.75;  $H_2O$ , 6.10 (desorption Karl-Fisher).

10 E.  $[3\alpha R-(3\alpha\alpha, 3''\alpha\alpha, 4\beta, 4''\beta, 5\alpha, 5''\alpha, 6\beta, 6''b, 7\alpha, 7''\alpha, 7\alpha\alpha, 7''\alpha\alpha)]$ -dodecahydro-4,4'',5,5'',6,6'',7,7''-octakis[[2-hydroxy-3-[(4,7,10-tricarboxymethyl)-1,4,7,10-tetraazacyclododecan-1-yl]propyl]oxy]dispiro-[1,3-benzodioxole-2, 1'-cyclo-hexane-4',2''-[1,3]benzodioxole],  
15 octagadolinium salt


Compound D (250 mg, 68.4  $\mu$ mol as the octatriethylammonium salt) in water (5 mL) at pH 6.0 was treated with tetrahydrated gadolinium acetate (222 mg, 546  $\mu$ mol) at 60°C for 5 hours. The pH of the aqueous  
20 reaction mixture was maintained at  $7.0 \pm 1.0$ . The resulting solution was then applied to a CHP-20P column (2.5 x 20 cm). The column was eluted with water (750 mL), 2.5% (300 mL), 5% (300 mL) and 10% (300 mL) of ethanol. The fractions containing the  
25 desired compound, which were eluted by 10% ethanol, were combined and concentrated in vacuo to obtain the octameric gadolinium chelate (200 mg) as a white solid in 60% yield.

Analysis Calculated for  $C_{154}H_{244}N_{32}O_{68}Gd_8 \cdot 19.58 H_2O \cdot 2.0 C_2H_5OH$ : C, 35.57; H, 5.58; N, 8.40. Found: C, 35.06; H, 6.22; N, 8.42;  $H_2O$ , 6.61% (desorption Karl-Fisher).

- 48 -

Example 7

(2S-(2 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ ,11 $\alpha$ ,)))-2,5,8,11-Tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid.  
 5 gadolinium salt

A. L-2-Benzylamino propanol

10 To a solution of N-benzoyl-L-alanine (29.0 g, 150 mmol) in tetrahydrofuran (200 ml) at 0°C was added a tetrahydrofuran solution of diborane (1 M solution, 800 ml) and the mixture was refluxed for 18 hours. Excess of diborane was decomposed with  
 15 methanol and the solvents were removed under reduced pressure. The residue was dissolved in methanol (100 ml) and treated with 6 N hydrochloric acid (100 ml). The mixture was heated at 70°C for 12 hours and the solvents were removed under reduced pressure. The  
 20 residue was co-evaporated with methanol (5 x 150 ml) and the product was dissolved in water (50 ml), basified with 5 N sodium hydroxide to pH 12 and extracted with ethyl acetate (3 x 200 ml). The combined organic layers were washed with saturated  
 25 sodium chloride (150 ml), dried and concentrated to a final volume of 100 ml. 300 ml of hexane was added and the solution cooled in the refrigerator

- 49 -

overnight. The white crystalline needles deposited were collected and dried to afford pure compound A (41.4 g, yield 87%). m.p. 46-48°C.

5 B. (S)-1-Benzyl-2-methyl-ethyleneimine

To a solution of Compound A (33.0 g, 200 mmol) and triphenyl phosphine (79.66 g, 300 mmol) in ether (500 ml) stirred under nitrogen in an ice bath, was slowly added diethyl azo-dicarboxylate (95%, 50 ml, 300 mmol). The solution was stirred at room temperature for 16 hours. A crystalline precipitate (triphenylphosphine/diethyl hydrazine carboxylate complex) was filtered off and washed with hexane/ether (1:1, 200 ml). The ether solution was extracted with 1 N hydrochloric (2 x 100 ml). The hydrochloric solution was basified with 5N sodium hydroxide, extracted with ether (3 x 150 ml), dried and concentrated to afford the crude aziridine compound B as a yellow oily liquid. This was further purified by distillation under reduced pressure to afford pure compound B as a colorless liquid (24.3 g, yield 75%). b.p. 71-72°C at 4 mm.

25 C. [2S-(2 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ ,11 $\alpha$ )]-2,5,8,11-Tetramethyl-1,4,7,10-tetrakis(phenylmethyl)-1,4,7,10-tetraazacyclododecane

To a solution of compound B (19.1 g, 130 mmol) in ethanol (250 ml) was added p-toluenesulfonic acid (PTSA, 1.1 g, 6.5 mmol) and the mixture was stirred at room temperature for 64 hours. At the end of this time, an additional 1.1 g (6.5 mmol) of PTSA was added and the mixture was stirred for 48 hours. Ethanol was removed in vacuo, and the product was purified by column chromatography over silica gel

- 50 -

(400 g) using methanol as the eluent. Fractions containing the pure product were combined, and solvent removal afforded 5.8 g of a salt. This was dissolved in methanol (100 mL) and basified with concentrated ammonia solution. The precipitated solid was filtered, dried and recrystallized from absolute ethanol to afford pure compound C as a colorless microcrystalline solid (2.8 g, yield 14%).  
m.p.: 147-148°C

10

D. [2S-(2 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ ,11 $\alpha$ ,)]-2,5,8,11-Tetramethyl-  
1,4,7,10-tetraazacyclododecane

To a solution of compound C (2.35 g, 4 mmol) in ethyl acetate (40 mL) and ethanol (400 mL) was added ammonium formate (2.52 g) and palladium acetate on Carbon (20%, 2.35 g). The mixture was stirred under reflux for 16 hours. The solution was filtered to remove the catalyst and the solvents were removed to afford pure Compound D as a light yellow solid (840 mg, yield 92%).

20

E. [2S-(2 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ ,11 $\alpha$ ,)]-2,5,8,11-Tetramethyl-  
1,4,7,10-tetraazacyclododecane-1,4,7,10-  
tetraacetic acid

To a solution of compound D (570 mg, 2.5 mmol) in acetonitrile (200 mL) was added potassium carbonate (4.0 g) and t-butyl bromo acetate (2.34 g, 12 mmol). The mixture was stirred at room temperature for 18 hours. Potassium carbonate was filtered off, acetonitrile removed in vacuo and the residue purified by column chromatography over silica gel (50 g) using chloroform and methanol to afford the tetra butyl ester (1.6 g). This material was dissolved in trifluoroacetic acid (150 mL), anisole

30

- 51 -

(10 mL) was added, and the mixture was stirred at room temperature for 16 hours. Trifluoroacetic acid was removed in vacuo and anisole was removed by co-evaporation with water (6 x 50 mL) to afford the crude product. The residue was dissolved in water (100 mL) and purified by anion exchange column chromatography over AG1-X2 resin (150 mL). The column, after washing with water, was eluted with 1 M formic acid. The fractions containing the pure product were combined. Solvent removal afforded the pure product. A small amount of formic acid that remained in the sample was removed by co-evaporation with water (5 x 50 mL) to obtain pure title compound as a colorless glassy solid (550 mg, yield 87%).

F. [2S-(2 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ ,11 $\alpha$ ,)]-2,5,8,11-Tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, gadolinium salt

A sample of Compound E (0.1 g) was dissolved in 8 mL deionized water. 0.04 mL 5N sodium hydroxide (NaOH) was added to convert the ligand into the monosodium salt (reactant I). In a separate vial, 0.08 g of GdCl<sub>3</sub> powder was dissolved in 1 mL deionized water (reactant II). At about 70°C and constant stirring, reactant II was added into reactant I dropwise over a time period of several hours. Meanwhile, 5N NaOH was pipetted into the solution from time to time to neutralize the protons released from chelation and maintain the reaction pH around 6. The excess metal ions were precipitated in the form of M(OH)<sub>3</sub>·xH<sub>2</sub>O by raising solution pH to about 9.5 and incubating at both about 70°C and room temperature for several hours. The precipitate was subsequently removed through centrifugation and

- 52 -

filtration with a 0.22  $\mu$ m membrane. The filtrate was finally condensed and neutralized to pH 7 in preparation for HPLC purification.


The crude chelate product was purified by preparative HPLC, using mobile phase gradient (CH<sub>3</sub>CN:H<sub>2</sub>O) and silica-based reversed phase column (YMC C18, 5 $\mu$ , 120 Å). The solvent was removed from fractions containing the desired product. The residue was dried in a vacuum oven at about 70°C overnight, furnishing the chelate in 65% yield. Mass Spectrum (FAB, m/e): Gd<sub>159</sub> + Na)<sup>+</sup> at 638. Elemental Analysis (C, H, N): Calculated for C<sub>20</sub>H<sub>33</sub>N<sub>4</sub>O<sub>8</sub>GdNa • 5.01 H<sub>2</sub>O: C, 33.00; H, 5.96; N, 7.70%. Found: C, 32.94; H, 5.63; N, 7.75%

15

- 53 -

Example 8

- [ $\alpha R-(\alpha R^*, \alpha' R^*, \alpha'' R^*, \alpha''' R^*, 2S^*, 5S^*, 8S^*, 11S^*)$ ]- $\alpha$ ,  
 $\alpha', \alpha'', \alpha'''$ , 2, 5, 8, 11-Octamethyl-1, 4, 7, 10-  
 5 tetraazacyclododecane-1, 4, 7, 10-tetraacetic acid,  
gadolinium salt



- 10 A. [ $\alpha R-(\alpha R^*, \alpha' R^*, \alpha'' R^*, \alpha''' R^*, 2S^*, 5S^*, 8S^*, 11S^*)$ ]- $\alpha$ ,  
 $\alpha', \alpha'', \alpha'''$ , 2, 5, 8, 11-Octamethyl-1, 4, 7, 10-  
 tetraazacyclododecane-1, 4, 7, 10-tetraacetic  
acid

- To a solution of compound D from Example 7  
 15 (570 mg, 2.5 mmol) in acetonitrile (200 ml) was added  
 potassium carbonate followed by L-benzyl-2-  
 trifluoromethanesulfonyloxy-propionate (prepared fresh  
 from benzyl lactate as described in S.I. Kang et al.,  
 Inorganic Chem., 1993, 32, 2912 -2918) (3.9 g, 12.5  
 20 mmol) and the mixture was stirred at room temperature  
 for 48 hours. Potassium carbonate was filtered off,  
 acetonitrile removed in vacuo and the residue  
 purified by column chromatography over silica gel  
 (100 g) using chloroform and the methanol to afford a  
 25 tetra benzyl ester (1.1 g). This material was  
 dissolved in a mixture of ethanol (75 ml) and water  
 (10 ml) and hydrogenated over 10% Pd/C (250 mg) for

- 54 -

18 hours. The catalyst was filtered off and solvent removal afforded the crude product. This was dissolved in water (100 ml) and purified by anion exchange column chromatography over AG1-X2 resin (150 ml). The column, after washing with water, was eluted with a gradient of 0-200 mM formic acid. Fractions containing the pure product were combined. Solvent removal afforded the pure product. A small amount of formic acid that remained in the sample was removed by co-evaporation with water (5 x 50 ml) to obtain pure Compound A as a colorless glassy solid (340 mg, yield 26%).

B.  $[\alpha R-(\alpha R^*, \alpha' R^*, \alpha'' R^*, \alpha''' R^*, 2S^*, 5S^*, 8S^*, 11S^*)] - \alpha, - \alpha', \alpha'', \alpha''', 2, 5, 8, 11$ -Octamethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, gadolinium salt

A sample of Compound A (0.15 g) was dissolved in 8 mL deionized water. 0.056 mL 5N NaOH was added to convert the ligand into the monosodium salt (reactant I). In a separate vial, 0.11 g of GdCl<sub>3</sub> powder was dissolved in 1 mL deionized water (reactant II). Reactants I and II were treated as described in the method of Example 7F to yield 74% of the title compound.

Mass Spectrum (FAB, m/e): (Gd<sup>159</sup> + Na)<sup>+</sup> at 694.

Elemental analysis: Calculated for

C<sub>24</sub>H<sub>40</sub>N<sub>4</sub>O<sub>8</sub>GdNa·1.98 H<sub>2</sub>O: C, 39.57, H, 6.08, N, 7.69%.

Found: C, 39.31, H, 6.19, N, 7.60%.




- 55 -

Example 9

[2S-(2R\*,5R\*,8R\*,11R\*)]-2,5,8,11,-Tetramethyl-  
1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid,  
gadolinium salt

5



A. [2S-(2R\*,5R\*,8R\*,11R\*)]-2,5,8,11-Tetramethyl-  
1,4,7,10-tetraazacyclododecane-1,4,7-triacetic  
acid

10

To a solution of Compound D (456 mg, 2 mmol) in acetonitrile (150 mL) was added sodium bicarbonate (4.0 g) and t-butyl bromo acetate (2.34 g, 12 mmol), and the mixture was stirred at room temperature for 48 hours. Sodium bicarbonate was filtered off, acetonitrile was removed *in vacuo* and the residue was dissolved in TFA (25 mL). Anisole (1 mL) was added and the mixture was stirred at room temperature for 12 hours. TFA was removed *in vacuo* and anisole was removed by co-evaporation with water (5 x 50 mL) to afford the crude product. The residue was dissolved in water (60 mL) and purified by anion exchange column chromatography over AG1-X2 resin (100 mL). The column, after washing with water, was eluted with a gradient of water to 50 mm M formic acid. The fractions were analyzed by HPLC and those containing the pure product were combined. Solvent removal afforded Compound A. A small amount of formic acid

15

20

25

- 56 -

that remained in the sample was removed by co-evaporation with water (5 x 50 mL) to obtain pure Compound E as a colorless glassy solid (320 mg, yield 40%).

5

B. [2S-(2R\*,5R\*,8R\*,11R\*)]-2,5,8,11,-Tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, gadolinium salt

10 A sample of Compound A (0.125 g) was dissolved in 9 mL deionized water (reactant I). In a separate vial, 0.129 g of GdCl<sub>3</sub> powder was dissolved in 1.5 mL deionized water (reactant II). Reactants I and II were treated as described in the method of Example 7F  
15 to yield 80% of the title compound.

Mass Spectrum (FAB, m/e): (Gd<sup>159</sup> + H)<sup>+</sup> at 559.

Elemental Analysis (C,H,N):

Calculated for C<sub>18</sub>H<sub>31</sub>N<sub>4</sub>O<sub>6</sub>Gd•2.81 H<sub>2</sub>O: C 35.60, H 6.08, N 9.22%.


20 Found: C 35.65, H 6.11, N 9.02%.

- 57 -

Example 10

10-(Phosphonomethyl)-1,4,7,10-tetraaza-  
cyclododecane-1,4,7-triacetic acid, gadolinium salt

5



A. 10-(Phosphonomethyl)-1,4,7,10-tetraazacyclo-  
dodecane-1,4,7-triacetic acid

10 A mixture of DO3A (1g, 2.89 mmol), phosphorus  
 acid (0.485 g, 5.91 mmol), concentrated hydrochloric  
 acid (1.1 mL, 13.4 mmol) and formaldehyde (1.35 mL of  
 37% aqueous solution, 16.4 mmol) in water (2.5 mL)  
 was refluxed for 27.5 hours. Removal of water from  
 15 the reaction mixture *in vacuo* gave an off-white solid  
 (1.74 g) as a crude product of Compound A. The solid  
 was dissolved in water (50 mL), and the pH of the  
 resulting solution was adjusted to 5.0 by addition of  
 1.0 N sodium hydroxide. The aqueous solution was  
 20 applied to a strong cation exchange column of AG50WX8  
 (250 mL, flowrate, 12.5 mL/minute). Initially, water  
 was used to remove any negatively charged inorganic  
 species, and ammonium hydroxide (1.0 M) was employed  
 to bring out the crude product. After removal of the  
 25 ammonium hydroxide from the fractions containing the  
 product, the residue dissolved in 5mM TEAB (pH 7.5)  
 was applied to a DEAE Sephadex ion exchange column  
 (750 mL, flowrate, 4 mL/minute). The column was

- 58 -

eluted with TEAB, whose concentration varied from 5 to 400 mM. The concentration of the buffer was doubled at every column volume (1L): 5, 10, 20, 40, 67, 80, 100, 125, 200 and 400 mM. Compound A  
5 required 400 mM TEAB to be brought out from the column. Evaporation of the buffer from the fractions containing Compound A gave a white residue (0.9 g). The solid, dissolved in water (35 mL) was loaded on a  
10 strongly basic anion column of Amberlite IRA 900 C (200 mL, flowrate, 14 mL/minute). Water was first used to remove triethylamine followed by sulfuric acid (1M) to elute Compound A from the column. The fractions containing the product were combined and applied to a column of Reillex 425  
15 poly(4-vinyl)pyridine (PVP) 250 mL, 10 mL/minute). Water was used to elute the product free from sulfuric acid. Removal of water from the eluate (3.7 L) yielded Compound A (0.6 g, 47%) as a dense solid.

20 B. 10-(Phosphonomethyl)-1,4,7,10-tetraaza  
cyclododecane-1,4,7-triacetic acid, gadolinium  
salt

A sample of Compound A (0.2 g) was dissolved in 5 mL deionized water and its pH was adjusted to 4  
25 with dilute NaOH. 0.096 g of Gd<sub>2</sub>O<sub>3</sub> powder was added slowly. The solution was refluxed overnight. The excess metal ions were precipitated in the form of M(OH)<sub>3</sub> · x H<sub>2</sub>O by raising solution pH to about 9.5 with 1N NaOH and incubating at both about 70°C and  
30 room temperature for several hours. The precipitate was subsequently removed through centrifugation and filtration with 0.22 µm membrane. The filtrate was finally condensed and neutralized to pH 7 with 1N HCl in preparation for HPLC purification. HPLC

- 59 -

purification was performed as in Example 7F to yield 64% of the title compound.

Mass Spectrum (FAB, m/e):  $(\text{Gd}^{159} + 2\text{Na-H})^+$  at 640.

Elemental Analysis (C,H,N): Calculated for


5  $\text{C}_{15}\text{H}_{24}\text{N}_4\text{O}_9\text{PGdNa}_2 \cdot 2.59 \text{ H}_2\text{O}$ : C 26.29, H 4.29, N 8.18%.

Found: C 26.23, H 4.18, N 7.93%.

- 60 -

What is claimed is:

1. A compound of the following formula I:



5

wherein

each m, n, o and p is independently 1 or 2;

q is 0 or 1;

10 each G is independently -COOR<sup>a</sup>, -P(O)(OR'')<sub>2</sub>,  
-P(O)(OR'')(R'') or -C(O)N(R'')<sub>2</sub>;

each R' is independently hydrogen or alkyl,  
alkoxy, cycloalkyl, hydroxyalkyl or aryl, each of  
which is optionally substituted, or a functional  
15 group capable of forming a conjugate with a  
biomolecule or of forming a multimer of said compound  
of formula I;

each R'' is hydrogen;

each R<sup>13</sup> through R<sup>20</sup> is independently  
20 hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl or a  
functional group capable of forming a conjugate with  
a biomolecule or of forming a multimer of said  
compound of formula I;

or R<sup>13</sup> together with R<sup>15</sup>, and R<sup>17</sup> together with  
25 R<sup>18</sup>, independently form, together with the carbon  
atoms in the tetraazacyclododecane macrocycle to  
which they are attached, a fused fully or partially

- 61 -

saturated non-aromatic cyclohexyl ring which may be unsubstituted or substituted by one or more halogen, alkyl, ether, hydroxy or hydroxyalkyl groups, and which may be further fused to a carbocyclic ring, or

5 R<sup>13</sup> and R<sup>15</sup>, are each hydrogen and R<sup>17</sup>, together with R<sup>18</sup>, forms a fused fully or partially saturated non-aromatic cyclohexyl ring as defined above, or R<sup>13</sup>, together with R<sup>15</sup>, forms a fused fully or partially saturated non-aromatic cyclohexyl ring as defined

10 above and R<sup>17</sup> and R<sup>18</sup> are hydrogen; provided that (a.) when G is always -COOR'' and (i.) R', R'', R<sup>14</sup> and R<sup>16</sup> through R<sup>20</sup> are all hydrogen, then R<sup>13</sup> and R<sup>15</sup> are other than hydrogen; (ii.) R'' and R<sup>13</sup> through R<sup>20</sup> are all hydrogen, and m, n, o, p and q are each 1,

15 then (CR'R') is other than (CH<sub>2</sub>) and (CHCH<sub>3</sub>); (iii.) R', R'', R<sup>13</sup>, R<sup>14</sup>, R<sup>17</sup> and R<sup>20</sup> are all hydrogen, then at least two of R<sup>15</sup>, R<sup>16</sup>, R<sup>18</sup> and R<sup>19</sup> are other than methyl; and (iv.) R'', R<sup>16</sup>, R<sup>19</sup> and R<sup>20</sup> are all hydrogen, and each (CR'R') is independently (CHR') or

20 (CH<sub>2</sub>CHR'), then R<sup>13</sup> and R<sup>15</sup>, and R<sup>17</sup> and R<sup>18</sup>, are other than a fused ring; and (b.) when G is always -P(O)(OR'')<sub>2</sub>, -P(O)(OR'')(R'') or -C(O)N(R'')<sub>2</sub>, then at least one R' or R<sup>13</sup> through R<sup>20</sup> is other than hydrogen;

25 or a salt or multimeric form thereof.

2. A metal chelate, comprising a compound of claim 1 complexed with a metal atom.


30 3. The chelate of claim 2, wherein the metal is selected from atoms having an atomic number of 21 to 29, 42 or 57 to 83.

- 62 -

4. The chelate of claim 3, wherein said metal is gadolinium.

5. A compound of the following formula II:

5



wherein

Q is a 4- to an 8-membered carbocyclic ring  
10 which may be fully or partially saturated;


t is an integer from 2 to 16;

each R group is independently hydrogen, -OH,  
-CH<sub>2</sub>-A, -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A or a functional group  
capable of forming a conjugate with a biomolecule,  
15 provided that at least two of the R groups are  
selected from -CH<sub>2</sub>-A or -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A; and

A is a moiety capable of chelating a metal  
atom.

20

6. A compound of Claim 5 wherein A is



each m, n, o and p is independently 1 or 2;



- 63 -

q is 0 or 1;

each G is independently -COOR'', -P(O)(OR'')<sub>2</sub>,  
-P(O)(OR'')(R'') or -C(O)N(R'')<sub>2</sub>;

each R' is independently hydrogen or alkyl,  
5 alkoxy, cycloalkyl, hydroxyalkyl or aryl, each of  
which is optionally substituted, or a functional  
group capable of forming a conjugate with a  
biomolecule or of forming a multimer;

each R'' is hydrogen;

10 each R<sup>13</sup> through R<sup>20</sup> is independently  
hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl or a  
functional group capable of forming a conjugate with  
a biomolecule or of forming a multimer of said  
compound of the formula I;

15 or R<sup>13</sup> together with R<sup>15</sup>, and R<sup>17</sup> together with  
R<sup>18</sup>, independently form, together with the carbon  
atoms in the tetraazacyclododecane macrocycle to  
which they are attached, a fused fully or partially  
saturated non-aromatic cyclohexyl ring which may be  
20 unsubstituted or substituted by one or more halogen,  
alkyl, ether, hydroxy or hydroxyalkyl groups, and  
which may be further fused to a carbocyclic ring, or  
R<sup>13</sup> and R<sup>15</sup> are each hydrogen and R<sup>17</sup>, together with  
R<sup>18</sup>, forms a fused fully or partially saturated non-  
25 aromatic cyclohexyl ring as defined above, or R<sup>13</sup>,  
together with R<sup>15</sup>, forms a fused fully or partially  
saturated non-aromatic cyclohexyl ring as defined  
above and R<sup>17</sup> and R<sup>18</sup> are hydrogen; provided that  
(a.) when G is always -COOR'' and (i.) R', R'', R<sup>14</sup>  
30 and R<sup>16</sup> through R<sup>20</sup> are all hydrogen, then R<sup>13</sup> and R<sup>15</sup>  
are other than hydrogen; (ii.) R'' and R<sup>13</sup> through R<sup>20</sup>  
are all hydrogen, and m, n, o, p and q are each 1,  
then (CR'R') is other than (CH<sub>2</sub>) and (CHCH<sub>3</sub>); (iii.)  
R', R'', R<sup>13</sup>, R<sup>14</sup>, R<sup>17</sup> and R<sup>20</sup> are all hydrogen, then


- 64 -

at least two of R<sup>15</sup>, R<sup>16</sup>, R<sup>18</sup> and R<sup>19</sup> are other than methyl; and (iv.) R'', R<sup>16</sup>, R<sup>19</sup> and R<sup>20</sup> are all hydrogen, and each (CR'R') is independently (CHR') or (CH<sub>2</sub>CHR'), then R<sup>13</sup> and R<sup>15</sup>, and R<sup>17</sup> and R<sup>18</sup>, are  
 5 other than a fused ring; and (b.) when G is always -P(O)(OR'')<sub>2</sub>, -P(O)(OR'')(R'') or -C(O)N(R'')<sub>2</sub>, then at least one R' or R<sup>13</sup> through R<sup>20</sup> is other than hydrogen;

or a salt or multimeric form thereof.

10


7. A compound of claim 5, wherein A is



15 each R' is independently hydrogen, alkyl, alkoxy, hydroxyalkyl, aryl, aralkyl or arylalkoxy; each R'' is hydrogen; and each n is 1 or 2.


20

8. A compound of claim 5 wherein



is

- 65 -




each  $R^1$  through  $R^{12}$  group is independently hydrogen, -OH, -CH<sub>2</sub>-A, -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A or a  
 5 functional group capable of forming a conjugate with a biomolecule;


at least two of  $R^1$  through  $R^{12}$  are selected from -CH<sub>2</sub>-A or -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A; and

$R^8$  and  $R^9$  taken together may additionally form  
 10 the group -O-[C(RR)]-O- where each R is independently hydrogen or alkyl, or  $R^8$  and  $R^9$  taken together may form


- 66 -



9. A compound of claim 6 wherein



is a compound of the formula



5


- 67 -

wherein


each  $R^1$  through  $R^{12}$  group is independently hydrogen, -OH, -CH<sub>2</sub>-A, -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A or a functional group capable of forming a conjugate with a biomolecule;

at least two of  $R^1$  through  $R^{12}$  are selected from -CH<sub>2</sub>-A or -OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-A; and

$R^8$  and  $R^9$  taken together may additionally form the group -O-[C(RR)]-O- where each R is independently hydrogen or alkyl, or  $R^8$  and  $R^9$  taken together may form




10. A compound of claim 7, wherein



15

is

- 68 -




each  $R^1$  through  $R^{12}$  group is independently  
 5 hydrogen,  $-OH$ ,  $-CH_2-A$ ,  $-OCH_2CH(OH)CH_2-A$  or a  
 functional group capable of forming a conjugate with  
 a biomolecule;

at least two of  $R^1$  through  $R^{12}$  are selected  
 from  $-CH_2-A$  or  $-OCH_2CH(OH)CH_2-A$ ; and

10  $R^8$  and  $R^9$  taken together may additionally form  
 the group  $-O-[C(RR)]-O-$  where each  $R$  is independently  
 hydrogen or alkyl, or  $R^8$  and  $R^9$  taken together may  
 form

- 69 -



11. A metal chelate, comprising a compound of claim 5 complexed with a metal atom.

5

12. A metal chelate, comprising a compound of claim 6 complexed with a metal atom.

13. A conjugate, comprising a compound of claim 5 conjugated with a biomolecule.

10

14. A metal chelate, comprising a conjugate of claim 13 complexed with a metal atom.

- 70 -

15. A method for diagnostic imaging,  
comprising the steps of administering to a host a  
compound of claim 5, which compound is complexed with  
a metal, and obtaining a diagnostic image of said  
5 host.

16. The method of claim 15, wherein said  
image is a magnetic resonance image.

10 17. A method for diagnostic imaging,  
comprising the steps of administering to a host a  
conjugate of claim 13, which conjugate is complexed  
with a metal, and obtaining a diagnostic image of  
said host.

15

18. A compound selected from the group  
consisting of:

(1 $\alpha$ ,2 $\alpha$ ,4 $\beta$ ,5 $\beta$ )-10,10'-[(1,2,4,5-Tetrahydroxy-  
20 1,4-cyclohexanediyl)bis(methylene)]bis[1,4,7,10-  
tetraazacyclododecane-1,4,7-triacetic acid];

(3 $\alpha$ ,4 $\alpha$ ,5 $\beta$ ,6 $\alpha$ ,7 $\beta$ ,7 $\alpha$ )-10,10'10'',10'''-[[Hexahydro-2,2-  
dimethyl-1,3-benzodioxol-4,5,6,7-tetrayl]tetra(oxy)-  
25 tetra(2-hydroxy-3,1-propanediyl)]tetrakis[1,4,7,10-  
tetraazacyclododecane-1,4,7-triacetic acid];

3,4,5,6-Tetra-O-[2-hydroxy-3-[4,7,10-  
tris(carboxymethyl)-1,4,7,10-tetraazacyclododecan-1-  
30 yl]propyl]-myo-inositol;



- 71 -

(1 $\alpha$ , 2 $\alpha$ , 3 $\alpha$ , 4 $\beta$ , 5 $\alpha$ , 6 $\beta$ )-10, 10'-[(2, 3, 5, 6-tetrahydroxy-1, 4-cyclohexanediyl)bis(2-hydroxy-3, 1-propanediyl)]-bis[1, 4, 7-10-tetraazacyclododecane-1, 4, 7-triacetic acid];

5

10-[[1, 4-Dihydroxy-2, 5-bis[2-hydroxy-3-[4, 7, 10-tris-(carboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl]-propoxy]-4-[[4, 7, 10-tris(carboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl]methyl]cyclohexyl]methyl]-1, 4, 7, 10-tetraazacyclododecane-1, 4, 7-triacetic acid;

10

[3aR-(3a $\alpha$ , 3\*a $\alpha$ , 4 $\beta$ , 4\* $\beta$ , 5 $\alpha$ , 5\*a $\alpha$ , 6 $\beta$ , 6\*b, 7 $\alpha$ , -7\*a $\alpha$ , 7a $\alpha$ , 7\*a $\alpha$ )]-dodecahydro-4, 4\*, 5, 5\*, 6, 6\*, 7, 7\*-octakis-[[2-hydroxy-3-[(4, 7, 10-tricarboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl]propyl]-oxy]dispiro-[1, 3-benzodioxole-2, 1'-cyclo-hexane-4', 2'-[1, 3]-benzodioxole];

15

[2S-(2 $\alpha$ , 5 $\alpha$ , 8 $\alpha$ , 11 $\alpha$ , )]-2, 5, 8, 11-Tetramethyl-1, 4, 7, 10-tetraazacyclododecane-1, 4, 7, 10-tetraacetic acid;

20

[ $\alpha$ R-(aR\*,  $\alpha$ 'R\*,  $\alpha$ \*R\*,  $\alpha$ '''R\*, 2S\*, 5S\*, 8S\*, 11S\*)]- $\alpha$ , - $\alpha$ ',  $\alpha$ \*, a''', 2, 5, 8, 11-Octamethyl-1, 4, 7, 10-tetraazacyclododecane-1, 4, 7, 10-tetraacetic acid;

25

[2S-(2R\*, 5R\*, 8R\*, 11R\*)]-2, 5, 8, 11, -Tetramethyl-1, 4, 7, 10-tetraazacyclododecane-1, 4, 7-triacetic acid; and

30 10-(Phosphonomethyl)-1, 4, 7, 10-tetraazacyclododecane-1, 4, 7-triacetic acid.

19. A compound selected from the group consisting of:

- 72 -

(1 $\alpha$ , 2 $\alpha$ , 4 $\beta$ , 5 $\beta$ )-10, 10'-[(1, 2, 4, 5-Tetrahydroxy-1, 4-cyclohexanediyl)bis(methylene)]bis[1, 4, 7, 10-tetraazacyclododecane-1, 4, 7-triacetic acid], digadolinium salt;

5

(3 $\alpha\alpha$ , 4 $\alpha$ , 5 $\beta$ , 6 $\alpha$ , 7 $\beta$ , 7 $\alpha\alpha$ )-10, 10'10'', 10'''-[[Hexahydro-2, 2-dimethyl-1, 3-benzodioxol-4, 5, 6, 7-tetrayl]tetra(oxy)-tetra(2-hydroxy-3, 1-propanediyl)]tetrakis[1, 4, 7, 10-tetraazacyclododecane-1, 4, 7-triacetic acid],

10 tetragadolinium salt;

3, 4, 5, 6-Tetra-O-[2-hydroxy-3-(4, 7, 10-tris(carboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl)propyl]-myo-inositol, tetragadolinium salt;

15

(1 $\alpha$ , 2 $\alpha$ , 3 $\alpha$ , 4 $\beta$ , 5 $\alpha$ , 6 $\beta$ )-10, 10'-[(2, 3, 5, 6-tetrahydroxy-1, 4-cyclohexanediyl)bis(2-hydroxy-3, 1-propanediyl)]-bis[1, 4, 7-10-tetraazacyclododecane-1, 4, 7-triacetic acid], digadolinium salt;

20

10-[[[1, 4-Dihydroxy-2, 5-bis[2-hydroxy-3-(4, 7, 10-tris(carboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl)-propoxy]-4-[[4, 7, 10-tris(carboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl)methyl]cyclohexyl)methyl]-

25 1, 4, 7, 10-tetraazacyclododecane-1, 4, 7-triacetic acid, tetragadolinium salt;

[3 $\alpha$ R-(3 $\alpha\alpha$ , 3'' $\alpha\alpha$ , 4 $\beta$ , 4'' $\beta$ , 5 $\alpha$ , 5'' $\alpha$ , 6 $\beta$ , 6'' $\beta$ , 7 $\alpha$ , -7'' $\alpha$ , 7 $\alpha\alpha$ , 7'' $\alpha\alpha$ )]-dodecahydro-4, 4'', 5, 5'', 6, 6'', 7, 7''-

30 octakis-[[2-hydroxy-3-[(4, 7, 10-tricarboxymethyl)-1, 4, 7, 10-tetraazacyclododecan-1-yl]propyl]-oxy]dispiro-[1, 3-benzodioxole-2, 1'-cyclo-hexane-4', 2''-[1, 3]-benzodioxole], octagadolinium salt;

- 73 -

[2S-(2 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ ,11 $\alpha$ ,)]-2,5,8,11-Tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, gadolinium salt;

5 [  $\alpha$ R-( $\alpha$ R\*, $\alpha'$ R\*, $\alpha''$ R\*, $\alpha'''$ R\*,2S\*,5S\*,8S\*,11S\*)]- $\alpha$ , $\alpha'$ , $\alpha''$ , $\alpha'''$ ,2,5,8,11-Octamethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, gadolinium salt;

10 [2S-(2R\*,5R\*,8R\*,11R\*)]-2,5,8,11,-Tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, gadolinium salt; and

10-(Phosphonomethyl)-1,4,7,10-tetraaza-  
15 cyclododecane-1,4,7-triacetic acid, gadolinium salt.

20. A chelate other than [1R-(1R\*,4R\*,7R\*,10R\*)]- $\alpha$ , $\alpha'$ , $\alpha''$ , $\alpha'''$ -tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic  
20 acid or [1R-(1R\*,4R\*,7R\*)]- $\alpha$ , $\alpha'$ , $\alpha''$ -trimethyl-1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid possessing a stability greater than  $10^{15}$  M<sup>-1</sup> and capable of exhibiting an immobilized relaxivity between about 60 and 200 mM<sup>-1</sup>s<sup>-1</sup>/metal atom.

25

21. A chelate of claim 20 capable of exhibiting an immobilized relaxivity between about 70 and 150 mM<sup>-1</sup>s<sup>-1</sup>/metal atom.

30

22. A chelate of claim 20 capable of exhibiting an immobilized relaxivity between about 80 and 100 mM<sup>-1</sup>s<sup>-1</sup>/metal atom.

- 74 -

23. A chelate possessing a stability greater than  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 1.

5

24. A chelate possessing a stability greater than  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 1.

10

25. A chelate possessing a stability greater than  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 1.

15

26. A chelate possessing a stability greater than  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 8.

20

27. A chelate possessing a stability greater than  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 8.

25

28. A chelate possessing a stability greater than  $10^{15} \text{ M}^{-1}$  and capable of exhibiting an immobilized relaxivity between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 8.

30

29. A chelate other than [1R-(1R\*,4R\*,7R\*,10R\*)]- $\alpha,\alpha',\alpha'',\alpha'''$ -tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid or [1R-(1R\*,4R\*,7R\*)]- $\alpha,\alpha',\alpha''$ -trimethyl-

- 75 -

1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid,  
including an immobilized functional group, capable of  
exhibiting an immobilized relaxivity between about 60  
and 200 mM<sup>-1</sup>s<sup>-1</sup>/metal atom.

5

30. A chelate of claim 29, capable of  
exhibiting an immobilized relaxivity between about 70  
and 150 mM<sup>-1</sup>s<sup>-1</sup>/metal atom.

10

31. A chelate of claim 29, capable of  
exhibiting an immobilized relaxivity between about 80  
and 100 mM<sup>-1</sup>s<sup>-1</sup>/metal atom.

15 32. A chelate including a functional group,  
capable of exhibiting an immobilized relaxivity  
between about 60 and 200 mM<sup>-1</sup>s<sup>-1</sup>/metal atom  
comprising a compound of claim 1.

20 33. A chelate including a functional group,  
capable of exhibiting an immobilized relaxivity  
between about 70 and 150 mM<sup>-1</sup>s<sup>-1</sup>/metal atom  
comprising a compound of claim 1.

25 34. A chelate including a functional group,  
capable of exhibiting an immobilized relaxivity  
between about 80 and 100 mM<sup>-1</sup>s<sup>-1</sup>/metal atom  
comprising a compound of claim 1.

30 35. A chelate including a functional group,  
capable of exhibiting an immobilized relaxivity  
between about 60 and 200 mM<sup>-1</sup>s<sup>-1</sup>/metal atom  
comprising a compound of claim 8.

- 76 -

36. A chelate including a functional group, capable of exhibiting an immobilized relaxivity between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 8.

5

37. A chelate including a functional group, capable of exhibiting an immobilized relaxivity between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 8.

10

38. A chelate other than [1R-(1R\*,4R\*,7R\*,10R\*)]- $\alpha,\alpha',\alpha'',\alpha'''$ -tetramethyl-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid, conjugated to a biomolecule, capable of exhibiting an immobilized relaxivity between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom.

15

39. A chelate of claim 38, capable of exhibiting an immobilized relaxivity between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom.

20

40. A chelate of claim 38, capable of exhibiting an immobilized relaxivity between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom.

25

41. A chelate conjugated to a biomolecule, capable of exhibiting an immobilized relaxivity between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 1.

30

42. A chelate conjugated to a biomolecule, capable of exhibiting an immobilized relaxivity between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom comprising a compound of claim 1.

- 77 -

43. A chelate conjugated to a biomolecule,  
capable of exhibiting an immobilized relaxivity  
between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom  
comprising a compound of claim 1.

5

44. A chelate conjugated to a biomolecule,  
capable of exhibiting an immobilized relaxivity  
between about 60 and 200  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom  
comprising a compound of claim 8.

10

45. A chelate conjugated to a biomolecule,  
capable of exhibiting an immobilized relaxivity  
between about 70 and 150  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom  
comprising a compound of claim 8.

15

46. A chelate conjugated to a biomolecule,  
capable of exhibiting an immobilized relaxivity  
between about 80 and 100  $\text{mM}^{-1}\text{s}^{-1}$ /metal atom  
comprising a compound of claim 8.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00337

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D257/02 C07D493/10 C07D405/14 A61K49/00 C07F9/6524  
 //(C07D493/10,317:00,317:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 382 583 (CELLTECH LIMITED) 16 August 1990 *Complete document* ---	1-4,6,7, 15-46
A	EP,A,0 365 412 (GUERBET S.A.) 25 April 1990 *Complete document* ---	1-4,6,7, 15-46
A	EP,A,0 485 045 (SCHERING AKTIENGESELLSCHAFT) 13 May 1992 *Complete document* *Page 28-29: example 21 ,sub b* -----	5,7-46



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*I\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

24 July 1995

Date of mailing of the international search report

27.07.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. ( + 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: ( + 31-70) 340-3016

Authorized officer

Luyten, H



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00337

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0382583	16-08-90	WO-A- 9009379 JP-T- 3504510	23-08-90 03-10-91
EP-A-0365412	25-04-90	FR-A- 2637895 AT-T- 122665 DE-D- 68922686 JP-A- 2196776 US-A- 5417960 US-A- 5049667	20-04-90 15-06-95 22-06-95 03-08-90 23-05-95 17-09-91
EP-A-0485045	13-05-92	DE-A- 4035760 AU-A- 8772691 CA-A- 2055093 JP-A- 4288063 US-A- 5277895	14-05-92 14-05-92 09-05-92 13-10-92 11-01-94